
HEALTH MANAGEMENT ASSOCIATES



Utilizing Zoom

- Ensure your audio is linked to your Zoom participant ID.
 - If you joined the audio by computer microphone and speaker, then you're all set.
 - If you joined the audio with a phone and did not enter your unique participant ID then enter # syour participant ID> # on your phone now. Note: Your unique participant ID can be found by clicking on the lower left corner of your Zoom screen where it soys Join Audio' and your Participant ID will appear.
- Ensure you are on MUTE and your camera is ON. On the bottom left corner of your screen, you will see a red line through the microphone.







Camera OFF

Microphone ON

Your participation throughout today via chat is appreciated!
Locate the chat box. On the bottom middle of your screen, click on the chat icon. This will open the "Zoom Group Chat" pane on the right side of your screen. You will see messages throughout the webinar on there. When prompted by the presenters, type in your answers or questions there.



HEALTH MANAGEMENT ASSOCIATES

2

Housekeeping

1

3

- Today is Session 1
- This series is eligible for both CEU's and CME's
 - These activities have been approved for CEU's by the Minnesota Board of Behavioral Health and Therapy for 3 hours of credit for LADCs and LPC/LPCCs (total of 12 hours if all four sessions are fully attended)
 - These activities have been approved for CME's by the American Academy of Family Physicians for 3 hours of credit (total of 12 hours if all four sessions are fully attended)
- · Please complete the evaluation for the webinar that will be sent out via email after each session.
- · You will be receiving a PDF of today's presentation.
- · This session is being recorded.

Follow-up questions? Contact Cami McIntire: cmcintire@healthmanagement.com

Welcome



Darin Rowles, MSW, LISW (he/him/his) Manager | HIV Supports, Disability Services Division Minnesota Department of Human Services

HEALTH MANAGEMENT ASSOCIATES

4

Acknowledgments

We would also like to thank our **community partners** for their support in developing this curriculum.











HEALTH MANAGEMENT ASSOCIATES

Land Acknowledgment



Every community owes its existence and vitality to generations from around the world who have contributed their hopes, dreams, and energy to making the history that led to this moment. Some were brought here against their will, some were drawn to leave their distant homes in hope of a better life, and some have lived on this land for more generations than can be counted. Truth and acknowledgment are critical to building mutual respect and connection across all barriers of heritage and difference.

We begin this effort to acknowledge what is buried by honoring the truth. We are standing on the ancestral lands of the Dakota people. We want to acknowledge the Dakota, the Ojibwe, the Ho Chunk, and the other nations of people who also call this place home. We pay respects to their elders past and present.

Please take a moment to consider the treaties made by the Tribal nations that entitle non-Native people to live and work on traditional Native lands. Consider the many legacies of violence, displacement, migration, and settlement that bring us together here today. Please join us in uncovering such truths at any and all public events.*

*This is the acknowledgment given in the USDAC Honor Native Land Guide – edited to reflect this space by Shannon Geshick, MTAG, Executive Director Minnesota Indian Affairs Council

HEALTH MANAGEMENT ASSOCIATES

5 6

Today's Presenters







Akiba Drew, MPH (she/her/hers) Associate Health Management Associates



Helen DuPlessis, MD, MPH (she/her/hers) Principal Health Management Associates

Disclosures

HEALTH MANAGEMENT ASSOCIATES

Faculty	Nature of Commercial Interest
Charles Robbins, MBA	Mr. Robbins discloses that he is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients.
Akiba Drew, MPH	Ms. Drew discloses that she is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients.
Helen DuPlessis, MD, MPH	Dr. DuPlessis discloses that she is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients. She is also a Board Member of Blue Shield of California Health Plan.
Jeanene Smith, MD	Dr. Smith discloses that she is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients.

HEALTH MANAGEMENT ASSOCIATES Cappright D 2021 Health Management Associates, Inc. All rights reserved. 7

8

Agenda for Webinar Series

Session	Topics
#1 WEDNESDAY, JAN 5 12:00 pm to 3:00 pm	Understanding HIV HIV Testing and Treatment The Science of Addiction Screening, and Assessment
#2 WEDNESDAY, JAN 12 12:00 pm to 3:00 pm	HIV Risk Reduction USUD Harm Reduction HIV and Stigma Motivational Interviewing Ethical and Legal Issues Funding and Policy Considerations
#3 WEDNESDAY, JAN 19 12:00 pm to 3:00 pm	Medications for Addiction Treatment Working with Justice Involved Persons Mental Health Treatment and Counseling Stimulant Use Chem Sex
#4 WEDNESDAY, JAN 26 12:00 pm to 3:00 pm	Cultural, Racial and Sexual Identities Pregnancy and HIV, SUD/OUD Accessing, Obtaining, and Integrating Services for Individuals with HIV and SUD in Minnesota

HEALTH MANAGEMENT ASSOCIATES

9

CHATTER FALL

Please respond to following prompt by typing into the chat box

> Please share a curiosity you bring with you today regarding the topics we are covering

Type your response and don't click enter.

HEALTH MANAGEMENT ASSOCIATES

10

SMALL BREAKOUT GROUPS

GET TO KNOW YOUR TRAINING COLLEAGUES

BREAKOUT ACTIVITY "Get to Know Your Colleagues"



INSTRUCTIONS

Step 1: Review How Breakouts Work

Step 2: Group Breakout 5 min

Share the following with the other participants in the room:

- Your personal pronouns
 Share one thing that you want people to know about you that relates this training

Step 3: Return to Main Room

BREAKOUT ACTIVITY "Small Breakout"





BREAKOUT ACTIVITY "Get to Know Your Neighbors"



INSTRUCTIONS

Step 1: Review How Breakouts Work

Step 2: Group Breakout 5 min Share the following with the other

- participants in the room: Name
- Your personal pronouns
- Share one thing that you want people to know about you that relates to this training

Step 3: Return to Main Room

13 14

BREAKOUT ACTIVITY "Get to Know Your Colleagues"



INSTRUCTIONS

Step 1: Review How Breakouts Work

Step 2: Group Breakout 5min

Share the following with the other participants in the room:

- Name
- Your personal pronouns
 Share one thing that you want people to know about you that relates to Austin


Step 3: Return to Main Room

Understanding HIV; HIV Testing and Treatment; The Science of Addiction, Screening and Assessment

Let's begin!

HEALTH MANAGEMENT ASSOCIATES

Pre-test Results



The Intersection of HIV and SUD

HEALTH MANAGEMENT ASSOCIATES

17

18

Context for the Intersection of HIV & SUD

- Substance use disorder (SUD) is frequently diagnosed among people with HIV.
- SUD also increases risk for acquiring HIV infection.
- The federal Health Resources and Service Administration (HRSA) recognizes the benefit of substance abuse treatment service for people with HIV and classifies outpatient treatment as a core medical service.

HEALTH MANAGEMENT ASSOCIATES

Context for the Intersection of HIV & SUD

- Tremendous biomedical advancements in HIV prevention and treatment have led to aspirational efforts to end the HIV epidemic.
- However, this goal will not be achieved without addressing the significant mental health and substance use problems among people living with HIV (PLWH) and people vulnerable to acquiring HIV.
- These problems exacerbate the many social and economic barriers to accessing adequate and sustained healthcare.

HEALTH MANAGEMENT ASSOCIATES

20

19

Glossary of Terms

- Sexual orientation a person's identity in relation to the gender or genders to which they are sexually
 attracted (straight, gay, lesbian, asexual, bisexual, pansexual)
- Gender identity and/or expression internal perception of one's gender; how one identifies or expresses oneself.
 - Cisgender a term used to describe a person whose gender identity aligns with those typically
 associated with the sex assigned to them at birth
 - Transgender refers to an individual whose current gender identity and/or expression differs from the sex they were assigned at birth (may have transitioned or be transitioning in how they are living)
 - Gender Expansive refers to an individual who expresses identity along the gender spectrum (genderqueer, gender nonconforming, nonbinary, agender, two spirit)
- Sexual Minority refers to a group whose sexual identity orientation or practices differ from the majority
 of and are marginalized by the surrounding society.

SOURCE: Centers for Educational Justice and Community Engagement, UC Berkele

- Race is usually associated with inherited physical, social and biological characteristics. In this context that
 means race is associated with biology. Institutionalized in a way that has profound consequences (White,
 African American, American Indian Alaskan Native, Native Hawaiian or Pacific Islander)*
- Ethnicity a term used to categorize a group of people with whom you share learned characteristics and identify according to common racial, national tribal, religious, linguistic, or cultural origin or background. (Hispanic, Non-Hispanic Black, Non-Hispanic Black, etc.)

SOURCE: US Office of Management and Budget: Federal Register Vol. 62(210): 58782

HEALTH MANAGEMENT ASSOCIATES

2021 Health Management Associates, Inc. All rights reserved.

22

Glossary of Terms

HEALTH MANAGEMENT ASSOCIATES

21

Common Acronyms

ART - Antiretroviral therapy

AUD – Alcohol use disorder

IDU - Injection or intravenous drug use

MAT – Medication assisted treatment or

Medications for addiction treatment

 $\ensuremath{\mathsf{MSM}}-\ensuremath{\mathsf{Men}}$ who have sex with men

OUD - Opioid use disorder

PEH - Person(s) experiencing homelessness

PEP - Post-exposure prophylaxis

PrEP – Pre-exposure prophylaxis

PLWH – Person(s) living with HIV

PWID – Person(s) who injects drugs SUD – Substance use disorder

HEALTH MANAGEMENT ASSOCIATES

Capyright 0 2221 Health Management Associates, Inc. As rights reserved.

Understanding HIV

Health Insurance Portability and Accountability Act (HIPAA) - required the creation of national standards to protect sensitive patient health information (PHI) from being disclosed without the patient's consent and includes a Privacy Rule addressing disclosure of and access to PHI; the Security Rule protects disclosure of and access to electronic PHI (e-PHI) a subset of information covered by the Privacy Rule

Code of Federal Regulations, Title 42, Part 2 (42 CFR Part 2) – a complicated set of regulations that strengthen the privacy protections afforded to persons receiving alcohol and substance use treatment (in addition to the more general privacy protections afforded in HIPAA). The regulations restrict the disclosure and use of alcohol

and drug patient records which are maintained in connection with any individual or entity that is federally assisted and holds itself out as providing, and provides, alcohol or drug abuse diagnosis, treatment or referral for treatment (42 CFR § 2.11)

Family Education Rights Protection Act (FERPA) - protects the privacy of student education records in public or private elementary, secondary, or post-secondary school and any state or local education agency that receives funds under an applicable program of the US Department of Education.

HEALTH MANAGEMENT ASSOCIATES

24

23

Learning Objectives:


HIV causes

illnesses



HEALTH MANAGEMENT ASSOCIATES

What is HIV?



HIV is the virus

Human: the virus can only infect human beings

Immunodeficiency: the virus destroys T-helper cells, an essential component of our body's immune system, leading to a deficiency in our body's ability to fight infection.

Virus: the organism is a virus which is incapable of reproducing by itself; it must use a human cell to

HEALTH MANAGEMENT ASSOCIATES

HIV is

transmitted

25

distinguish HIV and AIDS

26

What is HIV?



HEALTH MANAGEMENT ASSOCIATES

- · Ribonucleic acid (RNA) virus
- Classified as retrovirus (the virus inserts a copy of its genetic material (RNA) into the DNA of a host human
- Spread from person-to-person contact by contact with certain body fluids
- Weakens the immune system of a person by replicating inside T cells, a type of white cell also known as CD4 cells. The T cells are destroyed during this process.
- Once established, infection with HIV is chronic.
- · HIV is the virus that causes AIDS.

Time for a Poll



Approximately how many people in the United States are living with HIV?

- A. 275,000
- B. 500,000
- C. 1,100,000

D. 2,300,000

HEALTH MANAGEMENT ASSOCIATES

27

28

HIV Quick Facts



- · HIV is a chronic manageable infection
- Approximately 1.1 million people are living with HIV in the United States
- In 2019, an estimated 34,800 new HIV infections occurred in the United States.
- HIV continues to have a disproportionate impact on certain populations, particularly racial and ethnic minorities and gay, bisexual, and other men who have sex with men.

nttps://www.niv.gov/niv-basic

What is AIDS?

• AIDS is the disease:

HEALTH MANAGEMENT ASSOCIATES

- Acquired: HIV is not a condition passed on genetically; a person must become infected with it
- Immune: the immune system's ability to fight off viruses and bacteria becomes much less effective
- Deficiency: the immune system fails to work properly
- Syndrome: there are a wide range of diseases and opportunistic infections a person may experience once the immune system is depleted by HIV

HEALTH MANAGEMENT ASSOCIATES

Copyright © 2021 Health Management Associates, Inc. All rights reserved.

29

30

What is AIDS?

- It is a complex illness with a wide range of symptoms
- AIDS refers to individuals who have particular "AIDS—defining" disease such as:
 - a very low CD4 white blood cell count
 - specific illnesses acquired due to the weakened immune system (e.g., Burkitt's lymphoma, Kaposi sarcoma, pneumocystis pneumonia, toxoplasmosis, wasting syndrome)

Stages

1. Acute HIV infection

- HIV establishes infection in the body via replication within 11 days of initial acquisition
- · During acute infection, virus levels in the blood are very high.
- Very contagious
- Flu-like symptoms
- $^{\circ}$ $^{\circ}$ 50% of individuals will feel ill during acute infection

2. Chronic HIV infection

- Asymptomatic or latent
- Virus is active but is replicating at low levels
- · May last years
- Viral load increases, CD4 count decreases

3. AIDS


- CD4 < 200 cells/mm or opportunistic infections
- Can have high viral load and be infectious


Source: https://www.cdc.gov/hiv/basics/whatishiv.htm

HEALTH MANAGEMENT ASSOCIATES

HEALTH MANAGEMENT ASSOCIATES

31





Symptoms of HIV During Acute Infection

- Fevers
- Sore throat
- Chills
- Fatigue
- Rash
- Swollen lymph nodes
- Night sweats
- Mouth ulcers
- Muscle aches

HIV can not be diagnosed by symptoms, particularly those similar to other illnesses

HEALTH MANAGEMENT ASSOCIATES

Chronic Infection

34

- Once acquired, HIV is a lifelong infection
- There is no cure for HIV, but the infection can be controlled with medications much like diabetes.
- With treatment, the life expectancy of people with HIV is nearly the same as those who do not have
- Without treatment, most people living with HIV infection will go on to develop AIDS.

HEALTH MANAGEMENT ASSOCIATES

35 36

Latent HIV Reservoir

- Group of immune cells in the body that are infected with HIV but are not actively producing new HIV virus
- HIV medications do not affect these cells
- If a person stops taking their HIV medications, the infected cells in the reservoir can begin making new HIV virus

Source: https://hivinfo.nih.gov/understanding-hiv/fact-sheets/what-latent-hiv-reservoir

HEALTH MANAGEMENT ASSOCIATES

Copyright © 2021 Health Management Associates, Inc. All rights reserved.

37

CHATTER FALL

Please respond to following prompt by typing into the chat box

What information do you need to better prepare you to work with or care for individuals who are living with HIV?

Type your response and don't click enter.

HEALTH MANAGEMENT ASSOCIATES
Coordin't C 2021 Health Management Associates, Inc. All rights reserved.



38

References

UNDERSTANDING HIV

- "About HIV/AIDS." Centers for Disease Control and Prevention, https://www.cdc.gov/hiv/basics/whatishiv.html.
- "The Stages of HIV Infection." National Institutes of Health, U.S. Department of Health and Human Services, https://hivinfo.nih.gov/understanding-hiv/fact-sheets/stages-hiv-infection.



HEALTH MANAGEMENT ASSOCIATES

Copyright © 2021 Health Management Associates, Inc. All rights reserved.

HEALTH MANAGEMENT ASSOCIATES

40

39

HIV Incidence in Minnesota

- In 2020, there were 226 new HIV cases, which is a decrease from 276 cases in 2019
 - This is below the 5-year average of 274 cases
 - People assigned male gender at birth accounted for the majority of cases (84%)




- HIV rates among people who inject drugs is increasing:
 In 2014, 5% of new HIV cases were due to injection drug use

 - In 2019, 15% of new HIV cases were due to injection drug use
- Disparity
 - Over two-thirds (69%) of new cases are among communities of color despite these communities representing 17% of the

HEALTH MANAGEMENT ASSOCIATES

41

HIV Incidence in Minnesota




42

44

HIV Incidence in Minnesota

HIV Diagnoses# by County of Residence at Diagnosis, 2020




City of Minneapolis City of St. Paul Suburban* Greater Minnesota

59 cases (26%) 30 cases (13%) 84 cases (37%) 53 cases (24%)

HEALTH MANAGEMENT ASSOCIATES

HIV Incidence in Minnesota





HEALTH MANAGEMENT ASSOCIATES

43

HIV Prevalence in Minnesota

Estimated Number of people Living with HIV/AIDS in Minnesota

- As of December 31, 2020 9,422* people are assumed alive and living in Minnesota with HIV/AIDS. This includes:
 - 5,247 (56%) living with HIV infection (non-AIDS)
 - 4,175 (44%) living with AIDS
- This number includes 2,540 people who were first reported with HIV or AIDS elsewhere and subsequently moved to Minnesota
- This number excludes 1,629 people who were first reported with HIV or AIDS in Minnesota and subsequently moved out of state

*This number includes people with Minnesota reported as their current state of residence, regardless of residence at time of diagnosis. It also includes state prisoners and refugees arriving through the HIV+ Refugee Resettlement Program, as well as HIV+ Fredgees/immigrants arriving through other programs.

HEALTH MANAGEMENT ASSOCIATES

gyright © 2021 Health Management Associates, Inc. All rights reserved. 45

HIV Outbreak in Minnesota

HIV Outbreaks in Hennepin/Ramsey Counties & Duluth Region



Duluth area Outbreak
Declared 3/2021
Cases (N=21)*
Hennepin/Ramsey CountieOutbreak

Hennepin/Ramsey Counties Outbreak
Declared 2/2020
Cases (N=80)*
Disproportionate number of AI/AN

People who inject drugs People who are homeless or unstably housed

* Data counts as of 9/21/21

People at high-risk in the current outbreaks:

- People who inject drugs (PWID) or share needles/works
- People experiencing homelessness (PEH) or unstable housing
- People who exchange sex for income or other items they need

HEALTH MANAGEMENT ASSOCIATES

45

46

HIV Outbreak in Minnesota

- Synergistic with opioid epidemic
- Injection drug use is often a secondary effect of the over-prescription of opioids for pain as a core feature of the opioid epidemic

References

HIV IN MINNESOTA

- Minnesota Department of Human Services: HIV Resources https://mn.gov/dhs/people-we-serve/adults/health-care/hiv-aids/resources/
- "How the largest known homeless encampment in Minneapolis history came to be," The Appeal. July 15, 2020. https://theappeal.org/minneapolis-homelessness-crisis-powderhorn-park-encampment/
- "HIV Outbreak Response and Case Counts," Minnesota Department of Health. https://www.health.state.mn.us/diseases/hiv/stats/hiv.html
- "ACLU Minnesota, Mid-Minnesota Legal Aid file lawsuit to stop sweeps of homeless encampments," KARE 11. October 19, 2020. https://www.kare.ll.com/article/news/local/aclu-mn-files-suit-over-homeless-encampment-sweeps/39-3614595-3304-3602-3999-id-freebz/0653a
- "HIV/AIDS Statistics," Minnesota Department of Health. https://www.health.state.mn.us/diseases/hiv/stats/index.html
- "Health Advisory: HIV Outbreak and Syphilis Concern in Duluth Area," Minnesota Department of Health.
 March 4, 2021. https://www.health.state.mn.us/communities/ep/han/2021/mar4hiv.pdf
- "Health Advisory: HIV Outbreak in Persons Who Inject Drugs (PWID)," Minnesota Department of Health. February 6, 2020. https://www.health.state.mn.us/communities/ep/han/2020/feb3hiv.pdf
- "Quick Facts: Minnesota," U.S. Census. https://www.census.gov/quickfacts/MN

HEALTH MANAGEMENT ASSOCIATES

HEALTH MANAGEMENT ASSOCIATES

47



HIV Transmission

HIV is in:

- Blood
- Semen
- Vaginal fluids
- Anal fluids
- Breast milk

HIV is **not** in:

- Tears
- Sweat
- Insect bites
- Utensils
- Furniture, toilets

HEALTH MANAGEMENT ASSOCIATES

49

HEALTH MANAGEMENT ASSOCIATES

50

HIV Transmission

HEALTH MANAGEMENT ASSOCIATES

1. HIV must be present

a. One person must be currently infected with HIV

HIV Transmission (continued)

2. There needs to be enough virus

- a. Concentration of HIV determines whether infection will occur
- b. In the blood, the virus is very concentrated
 - i. Therefore, it can take a small amount of blood to infect someone
- c. In bodily fluids like semen, vaginal and anal fluids, or breastmilk, virus levels can change overtime
 - Therefore, the chances of transmitting HIV may be lower for those with lower viral loads

HEALTH MANAGEMENT ASSOCIATES

51

HIV Transmission (continued)

3. HIV must get into the bloodstream

- a. Infectious fluids:

 - Semen
 - Vaginal secretions Anal fluids
 - Breast milk
- b. HIV can enter through:

 - Open cut or sore
 Mucous membranes like the genitals, anus, and rectum

 - · HIV cannot cross healthy, unbroken skin
- c. Main transmission routes for the HIV virus:
 - Unprotected sexual intercourse
 - Sharing needles for injection drug use
 - · Mother to child transmission

HEALTH MANAGEMENT ASSOCIATES

Sexual Transmission

- · Most common HIV transmission route
- Presence of other sexually transmitted infections can increase the risk of HIV transmission
- Vaginal Sex
 - The female is at the greatest risk because the lining of the vagina is a mucous membrane which can provide easy access to the bloodstream for HIV carried in semen
- Anal Sex
 - · Without a condom, riskiest sexual activity for HIV
 - Receptive partner is at greatest risk
 - Cell wall of the rectum is very thin
 - Anal tissue can be easily bruised or torn during sex which then provides easy access to the bloodstream for HIV carried in semen
 - Insertive partner also at some risk because the membranes inside the urethra can provide entry for HIV into the bloodstream

HEALTH MANAGEMENT ASSOCIATES

53

54

Sexual Transmission (continued)

- Oral to Anal
 - · Poses minimal HIV risk
- - Mouth is an unfriendly environment for HIV
 - Saliva contains enzymes that break down the virus and the mucous membranes in the mouth are more protective than anal or vaginal tissue
 - There are a few documented cases where it appears that HIV was transmitted orally, and those cases are attributed to ejaculation into the mouth
 - · Risk only for the person performing the oral sex
 - With a female partner performing oral sex on a woman who is menstruating increases the risk because blood has more HIV than vaginal fluid

Time for a Poll



Approximately how many days can HIV survive in a syringe at room temperature?

- A. None
- B. 24 hours
- 5 days D. 10 days
- E. 21 days
- 42 days

HEALTH MANAGEMENT ASSOCIATES

HEALTH MANAGEMENT ASSOCIATES

56

55

Non-Sexual Transmission

- Typically involve medical settings or accident scenes where there is a very large volume of blood exposure or a needle stick
- Injection drug use

 - Very high risk for HIV transmission
 Sharing a syringe is the most efficient way as it passes blood directly from one person's blood stream to another's
 At room temperature, HIV can live as long as 21 days in a syringe

 - When the temperature is **cold** (near freezing), HIV can live up to **42 days in a syringe**
 - An HIV-negative person has a **1 in 160** chance for getting HIV every time they use a needle that has been used by someone with HIV.
- Tattoos and piercings

 - No documented cases
 But theoretical risk of transmission
- Mother to infant

 - By exposure to blood and vaginal fluids
 During birth or through breast milk during feeding

HEALTH MANAGEMENT ASSOCIATES

How Does HIV cause Illness?

- HIV reproduces continuously in the body from the first day of infection.
- Initial Stage:
 - · Individuals can experience severe flu-like symptoms
 - Initial stage can last 2-4 weeks
 - Immune system attacks HIV and can clear large amounts of the virus every 24 hours
 For each virus particle cleared, a new one is created.
 - Anti-HIV response temporarily created equilibrium between immune cells and HIV virus
 - Equilibrium can last for months or years
- After initial stage:

 - No outward signs of illness that can last for years
 HIV viral load increases and CD4 T cell count declines
 - · Immune system starts working improperly
 - HIV overwhelms immune system, leaving the body vulnerable to other illness-causing infections

HEALTH MANAGEMENT ASSOCIATES

57

58

HIV and Hepatitis C (HCV) Co-Infections

- HCV is a bloodborne virus transmitted through direct contact with the blood of an infected person.
- Co-infection is common (50%-90%) among HIV-infected injection drug users (CDC, 2014).
- In co-infected persons, age at time of HCV infection, immune cell (CD4) count and level of alcohol consumption are associated with a higher rate of liver fibrosis.
- Risk of HCV similar to those of HIV:
 - Transfusion prior to 1992
 - Injecting drug use (most common) · Long term hemodialysis

 - High risk sexual contact
 - Occupational exposures to blood or blood products
 - Receiving an organ or tissue transplant from someone infected with HCV
 - · Transmission from HCV-infected mother to infant.

HEALTH MANAGEMENT ASSOCIATES

HIV and Hepatitis C Co-Infections



HEALTH MANAGEMENT ASSOCIATES

60

59

HIV and Hepatitis C Co-Infections

- In 2018 in Minnesota, there were 60 acute HCV cases and 33,856 chronic cases
 - 8,140 Co-infected for HIV and HCV

61

 The U.S. Public Health Service/Infectious Diseases Society of America guidelines recommend that all HIVinfected persons be screened for HCV infection (CDC, 2014). References

HIV TRANSMISSION

- "HIV and Injection Drug Use", Centers for Disease Control and Prevention (2021), https://www.cdc.gov/hiv/basics/hiv-transmission/injection-drug-use.html.
- "How Is HIV Transmitted?", HIV.govDate (2019), https://www.hiv.gov/hiv-basics/overview/about-hiv-and-aids/how-is-hiv-transmitted.

HEALTH MANAGEMENT ASSOCIATES Capyright © 2021 Health Management Association, Inc. All rights reserved. 61

HEALTH MANAGEMENT ASSOCIATES

62



Learning Objectives:



Explain HIV testing and methods, including testing and treatment policies in MN; Describe ART, what does ART stand for, and how

it is used

relationship between substance use practices and increased risk of acquiring HIV Explain the relationships between HIV and Hepatitis C Describe the options and indications for preand postexposure prophylaxis (PrEP and PEP) and treatment of HIV infection

HEALTH MANAGEMENT ASSOCIATES

63

HEALTH MANAGEMENT ASSOCIATES

HIV Quick Facts



- Fewer than 40% of people in the United States have ever had an HIV test.
- Nationally, less than 30% of people in the United States most at risk of acquiring HIV were tested in the past year (gay, bisexual and other MSM, transgender women, and
- In the 50 local jurisdictions where more than half of HIV diagnoses occur, less than 35% of people recommended for annual HIV testing were tested in the past year.

HEALTH MANAGEMENT ASSOCIATES

65

Group Discussion

What myths or barriers exist that prevent more people from getting an HIV test?

Use the "raise your hand" feature in Zoom or simply come off mute.



HEALTH MANAGEMENT ASSOCIATES

66

HIV Testing

- First step in HIV diagnosis and preventing the spread of HIV
- Testing is a crucial step in engaging people living with HIV into care
- CDC recommends everyone 13 to 64 years old get tested for HIV at least once as part of their routine care
- Additionally, clients should be tested if the client:
 - Has engaged in risky behaviors
 - Has ever had a sexually transmitted infections (STI)
 - Has a history of sharing drug injection equipment
 - Is presenting with any of a number of symptoms that might indicate recent infection with HIV or early symptomatic infection

Type of Tests



- Testing has become more sophisticated over time more sophisticated tests (i.e., $4^{\rm th}$ or $5^{\rm th}$ generation tests) look for both HIV antibodies and antigens
- Antibody tests look for they body's antibodies to HIV in the blood or oral fluids
 - Measure immune response to HIV
 Not useful in acute infections

 - Rapid tests and FDA-approved HIV self tests
- Antigen tests detect actual particles of the HIV virus that trigger the body to make antibodies
- Antibody/Antigen tests detect both and most common test in the US
- Nucleic acid test (NAT) looks for the actual virus in blood. Very expensive. Can detect HIV infection 10 to 33 days after an exposure.

HEALTH MANAGEMENT ASSOCIATES

HEALTH MANAGEMENT ASSOCIATES

67

Rapid HIV Tests

- · Several FDA approved tests are available for use
- · Provides results in 10 to 40 minutes
- · Look for the presence of HIV antibodies
- · Either negative or reactive
 - · Negative means no HIV antibodies were detected

 - If individual has had three or more months without an HIV risk exposure, the person can be considered negative
 If individual has had exposure, the person should be tested again after three full months
 - · Reactive means antibodies have been detected
 - A confirmatory test is required before diagnosis is given
 - A Western Blot test is generally used as the confirmatory test

 - This is done with a blood draw and processed at a medical lab
 Results given in one to two weeks
 Michigan Department of Health (MDH) allows funded programs to do rapid to rapid confirmatory testing shortening this window
 - Can also use a more recent 4th generation antibody/antigen test to confirm

HEALTH MANAGEMENT ASSOCIATES



Minnesota Reporting

- In Minnesota, anonymous testing is no longer offered due to reporting requirements.
 - · Confidential testing continues to be available.
- Minnesota's reporting law requires testing sites to pass along all identifying information about the client to the Minnesota Department of Health (MDH).
- This means a testing client's information is only used if a test is reactive, and then only to facilitate the process of linking clients to care.
- · Getting clients into care soon after they test HIVpositive will greatly improve their health and decrease their chance of spreading the virus.

HEALTH MANAGEMENT ASSOCIATES

69

70

Treatment



- There are now many medications a person living with HIV can take to slow the progression of the disease.
- When taken as prescribed, these medications can keep a person's health stable for a very long time
- When taken as prescribed these medications can also greatly reduce the ability to pass HIV to others.

HEALTH MANAGEMENT ASSOCIATES

What Happens if Diagnosed HIV Positive?

- · A thorough medical history is an important step to help the clinician proceed to clinical evaluation and formulate a treatment plan.
- Before starting antiretroviral therapy (ART) in any patient, laboratory studies should be done and may include HIV ribonucleic acid (RNA) (or viral load), CD4+ T cell counts, blood counts, screening chemistries, syphilis, toxoplasmosis, purified protein derivative (PPD), hepatitis A, B, and C viruses, and chest x-ray.
- · All patients with HIV should be tested and begin treatment with antiretrovirals as soon as possible, regardless of disease status.
- Adherence should be maintained because non-adherence can lead to the rapid development of drug resistance and disease progression.
- · One means to encourage adherence is to educate clients and their significant others about HIV/AIDS treatment (TIP 37; SAMHSA, 2008).
- · It is difficult for unhoused individuals to maintain adherence

HEALTH MANAGEMENT ASSOCIATES

72

71

What is Antiretroviral Therapy?

- Antiretroviral therapy (ART)
 - · Medicines used to treat HIV
 - Do not cure or remove virus from the body
 - Stops the virus from replicating
 - · Combination of HIV medications taken daily
 - · From different drug classes
 - · Blocks HIV at different stages of HIV life cycle
 - · Goal: undetectable viral loads

"Undetectable" is now commonly defined as having fewer than 20 copies/mL because a lot of lab tests can now "detect" HIV at that level.

"Viral load suppression" is usually defined as having fewer than 200 copies of HIV per milliliter of blood (copies/mL).

HEALTH MANAGEMENT ASSOCIATES

73

Undetectable = Untransmissible (U=U)

- People cannot transmit the HIV through sexual contract when their viral load is undetectable
- Undetectable means too low to be measured (<200 copies per mL)
- This can take up to 6 months after initiating HIV medications
 - Confirmed by a blood test given by your doctor
 - Should be followed up with another blood test 6 months afterwards



HEALTH MANAGEMENT ASSOCIATES

74

U = U for Non-Sexual Transmission

- Undetectable viral loads also crucial to pregnancy, breastfeeding, and injection drug use
 - The risk of transmitting HIV during pregnancy with an undetectable viral load is one in one thousand
 - The risk is not eliminated during breastfeeding, but an undetectable viral load reduces the risk of passing HIV
 - Unsure of how much the risk is reduced when sharing needles during injection drug use



HEALTH MANAGEMENT ASSOCIATES

U = U and Sexual Partners

- Involving partners in treatment plan can help patients adhere to treatment
- Encourage HIV-positive patients to talk to current and potential partners about what undetectable means
- Counsel patients and their partners to use strategies to maintain healthy sexual lives
 - Condoms to prevent pregnancy and sexually transmitted infections (STIs)
 - HIV treatment adherence (ART) for an HIV-positive patient
 - Pre-exposure Prophylaxis (PrEP) and Post-exposure Prophylaxis (PEP) for an HIV-negative partner

HEALTH MANAGEMENT ASSOCIATES

75

Medication Resistance



- Stopping and re-starting treatment can cause drug resistance to develop
- People receiving intermittent ART have twice the rate of disease progression compared to those receiving continual treatment
- Transient increases in viral load followed by a dip back to undetectable called 'blips'
 - Blips are common and are not indicative of a treatment failure
- U.S. HIV treatment guidelines recommends viral load be measured every 3 4 months until undetectable, then less frequent

HEALTH MANAGEMENT ASSOCIATES

References

HIV TESTING AND TREATMENT

- Centers for Disease Control and Prevention (2019), "CDC Press Release: Most Americans Have Never Had an HIV Test, New Data Show." https://www.cdc.gov/media/releases/2019/p0627-americans-hiv-test.html.
- Minnesota Dept. of Health, "Undetectable = Untransmittable (U=U).", https://www.health.state.mn.us/diseases/hiv/prevention/uu/index.html.
- NYC Health, "HIV: Undetectable Equals Untransmittable (U=U)." https://www1.nyc.gov/site/doh/health/health-topics/hiv-u-u.page

HEALTH MANAGEMENT ASSOCIATES


77 78



HIV Prevention



- Safer sex practices like condom use
- · Antiretroviral advances
 - Can reduce HIV viral load to undetectable levels making it less likely to be transmitted
- Post-exposure Prophylaxis (PEP)
 - · For individuals who have been exposed to



• Reduce the risk of being infected with HIV by 92%-99%

HEALTH MANAGEMENT ASSOCIATES

20

79

HEALTH MANAGEMENT ASSOCIATES

PrEP

PrEP (pre-exposure prophylaxis) is medicine people at risk for HIV take to prevent getting HIV from sex or injection drug use. When taken as prescribed, PrEP is highly effective for preventing HIV.

- CDC endorsed PrEP for HIV prevention in May 2014
- · Once-daily pill
- · Taken by individuals at high risk including, but not limited to:
 - People who inject drugs
 - People with HIV+ sexual partners
 - Individuals who intermittently or never use condoms




HEALTH MANAGEMENT ASSOCIATES

PrEP and Women



DOMINATE



to prevent HIV • Does not require negotiation or disclosure

• Woman-controlled option

- Especially important for women experiencing intimate partner violence
- Yet, underutilized in women due to systemic barriers to access

HEALTH MANAGEMENT ASSOCIATES

81

82

PEP

PEP (post-exposure prophylaxis) means taking medicine to prevent HIV after a possible exposure.

PEP should be used only in emergency situations and must be started within 72 hours after a recent possible exposure to HIV.

- Taking medicine to prevent HIV after a possible exposure

 - Through needle sharing
 - · Occupational exposures such as needle sticks
 - If sexually assaulted
- Only used in emergency situations
- Two antiretroviral medications taken daily for 28 days
- · Afterwards, you need to return to doctor for a HIV test
- If you have frequent exposures to HIV, then PEP is not right for you. You should take PrEP.

HEALTH MANAGEMENT ASSOCIATES

HIV and COVID-19

- People with HIV may be more likely to get severely ill from COVID-19
- However, evidence suggests those virally suppressed are at no greater risk booster is still generally recommended but should be at the advice of their physician
- Vaccines are safe for HIV-positive patient
 - A third dose of mRNA COVID-19 vaccination is recommended after the initial two doses
 - · Booster shots are already available
- · However:

84

- · It may not fully protect them
- They should follow all precautions of an unvaccinated person
- They should continue taking their ART (or PrEP for uninfected individuals)
 - Make sure you have a 30- to 90-day supply of medicine, if possible

HEALTH MANAGEMENT ASSOCIATES

83

Impact of COVID-19 on Care

- \bullet Some STIs and HIV rates decreased during the pandemic by approximately 2%
- Disruptions to care and testing likely impacted the number of cases reported
 Disruptions to preventative care means fewer testing opportunities
- Presents challenges when tracking the two outbreaks currently happening



HEALTH MANAGEMENT ASSOCIATES

References

HIV PREVENTION

- CDC: Prevention Basics https://www.cdc.gov/hiv/basics/prevention.html
- NIH.GOV: The Basics of HIV Prevention: https://www.cdc.gov/hiv/basics/prevention.html

HEALTH MANAGEMENT ASSOCIATES

86

85



CHATTER FALL

Please respond to following prompt by typing into the chat box

> Please share a curiosity you bring with you today about the science, screening and/or assessment of SUD

Type your response and don't click enter.

HEALTH MANAGEMENT ASSOCIATES

87



The Science of Addiction, Screening, and Assessment

HEALTH MANAGEMENT ASSOCIATES

89 90

Learning Objectives:



Describe at least two ways in which dopamine influences OUD recovery and treatment

Explain the neurobiological contributions to developing and sustaining addiction Define and distinguish screening, assessment, and American Society of Addiction Medicine (ASAM) level of care determination Identify and explain the complex interactions between HIV and SUD, such as virial load, treatment retention and compliance and retroviral resistance

HEALTH MANAGEMENT ASSOCIATES

91

Time for a Poll




Which of the following do you think is the *root* cause of substance use disorders?

- a) Personal choice and behaviors
- b) Impact of trauma and other adverse life events
- c) Abnormalities of neurochemicals in the brain
- d) I haven't decided yet

HEALTH MANAGEMENT ASSOCIATES

92


Science of Addiction



HEALTH MANAGEMENT ASSOCIATES Cappright © 2021 Health Management Associates, Inc. All rights reserve


93

Science of Addiction




94

Comparative Dopamine Production




95

Science of Addiction



Substances Affect on the Brain



DSM-5: Diagnosis of Opioid Use Disorder (OUD)

Category	Criteria	
Impaired control	Opioids used in larger amounts or for longer than intended Unsuccessful efforts or desire to cut back or control opioid use Excessive amount of time spent obtaining, using, or recovering from opioids Craving to use opioids	
Social impairment	Failure to fulfill major role obligations at work, school, or home as a result of recurrent opioid use Persistent or recurrent social or interpersonal problems that are exacerbated by opioids or continued use of opioids despite these problems Reduced or given up important social, occupational, or recreational activitie because of opioid use	
Risky use	Opioid use in physically hazardous situations Continued opioid use despite knowledge of persistent physical or psychological problem that is likely caused by opioid use	
Pharmacological properties	Tolerance as demonstrated by increased amounts of opioids needed to achieve desired effect; diminished effect with continued use of the same amount Withdrawal as demonstrated by symptoms of opioid withdrawal syndrome; opioids taken to relieve or avoid withdrawal	


98

Dopamine Depletion effects Recovery: It takes time for your brain to recover

- + Prolonged drug use changes the brain in long lasting ways
- + Changes are both functional and structural
- + Return to normal dopamine production is under study (takes over 1 year)
- + Discontinuing treatment before brain recovery may affect outcomes




Dopamine Depletion Effects Recovery



100

25

Science of Addiction: Treatments



HEALTH MANAGEMENT ASSOCIATES

Time for a Poll



Which statement about screening & testing for SUD is the most accurate?

- A. Universal toxicology testing is the most equitable way to identify substance use disorder (SUD) across Minnesota.
- B. For some populations, screening for SUD using an evidence-based verbal screening tools is about as sensitive as using toxicology testing in identifying SUD.
- C. Urine and serum toxicology tests are so sensitive, their results don't require a confirmatory test.
- D. Hospitals can obtain a toxicology sample without obtaining consent.
- E. Decisions about what screening tools to use are generally made based on data from research studies.

HEALTH MANAGEMENT ASSOCIATES

101

102

Screening, Assessment, Level of Care

+ Screening:

A rapid evaluation to determine the possible presence (risk) of a condition (high sensitivity, usually low specificity)

A more detailed evaluation meant to solidify the presence of a disease and sometimes assess disease severity (lower sensitivity, high specificity)

+ Level of Care Determination:

Evaluation of various biopsychosocial and other factors to determine/recommend the most appropriate level of care for the severity of the condition identified (outpatient vs inpatient).

HEALTH MANAGEMENT ASSOCIATES

Screening

WHEN TO SCREEN?

Key is to screen patients to determine who should have further assessment


Times not to screen:

- + Recent screen → Set interval for repeat screening
- + Current/recent diagnosis of SUD
- + Presumptive positive
 - Legal involvement (substance related arrest, DUI)
 - Toxicology results
 - Patient report

HEALTH MANAGEMENT ASSOCIATES

103

Screening



Screening

VALIDATED SCREENING TOOLS

- + Screening tools are validated for use in specific populations
- + Screening for co-morbid conditions and suicide is also critical

General Population	Pregnant Women	Youth
National Institute for Drug Addiction (NIDA) – Quick Screen Tobacco, Alcohol, Prescription, and other Substances (TAPS) AUDIT (Alcohol only) Patient History Questionnaire (PHQ-9) General Anxiety Disorder (GAD-7) PTSD Checklist (PCL-5) Columbia Suicide Severity Rating Scale (C-CCRS)	+ NIDA – Quick Screen* + P's plus (license fee) + Substance Use Risk Profile – Pregnancy (SURP) + CRAFFT – for 12 - 26 yo women (Car, Relax, Alone, Forget, Friend/Family, Trouble) + Perinatal Mood and Anxiety Disorder (PMAD) – Edinburgh, PHQ-9	Brief Screener for Alcohol, Tobacco and other Drugs (BSTAD) (12-17yo) Screening to Brief Intervention (S2BI) (12- 17yo) Problem oriented screening instrument for Teens (POSIT) CRAFFT*

HEALTH MANAGEMENT ASSOCIATES

105 106

Screening

A BRIEF WORD ABOUT TOXICOLOGY TESTING TERMINOLOGY

- + <u>Screen</u>: a qualitative (detected/ not detected) test; usually designed to detect many drug classes; confidence in results may be poor but depends on the assay. Also called preliminary immunoassay point of care test (POC).
 - + Make sure you know what is covered by your toxicology panel
- Confirmation: a test designed for very high confidence in identification of individual drugs/compounds; may be qualitative or quantitative (reports the amount of drug present).
- <u>Cutoff</u>: the concentration above which the substances is indicated as detected & below which the result indicates the substance was not detected; defined by the "kit" manufacturer, or by the limit of quantification (LOQ).
 - + Knowing your lab cutoff values can avoid action on false positives (e.g., poppy seeds, oxycodone and hydrocodone)

Screening: Is There a Role of Toxicology Testing?

- Typically does not test for alcohol or tobacco use
- Potential for false positive and false negative results
- Complicated relationship between toxicology and child welfare involvement
- Test results do not assess parenting capabilities or other qualities
- Often applied selectively
- · Lab cut-off points for sensitivity
- Positive toxicology test does not establish the diagnosis of SUD



HEALTH MANAGEMENT ASSOCIATES

Capyone 2 2021 health theorement homeone, inc. As a option recovered against the properties of the control of t

107 108

Screening

BEST PRACTICES FOR SCREENING: USE MOTIVATIONAL INTERVIEWING TO START A CONVERSATION

+ "An important part of primary care/prenatal care [supporting you to stay with / reclaim custody of your baby] is screening for any risky conditions. Some of these conditions can be scary to talk about but are pretty common. Also, no matter the issue we have the ability to help work through it."

Is it ok if I ask you some questions about those risks?

+ For someone in treatment... We're doing a urine drug test today, will there be any findings on that test I'm not expecting?

HEALTH MANAGEMENT ASSOCIATES

109

Poll Answer

WHICH STATEMENT ABOUT SCREENING & TESTING FOR SUD IS THE MOST ACCURATE?

- A. Universal toxicology testing is the most equitable way to identify substance use disorder (SUD) across Minnesota.
- B. For some populations, screening for SUD using an evidence-based verbal screening tools is about as sensitive as using toxicology testing in identifying SUD.
- C. Urine and serum toxicology tests are so sensitive, their results don't require a confirmatory test.
- D. Hospitals can obtain a toxicology sample without obtaining consent.
- E. Decisions about what screening tools to use are generally made based on data from research studies.

HEALTH MANAGEMENT ASSOCIATES

110

S(A)BIRT

SCREENING BRIEF INTERVENTION AND REFERRAL TO TREATMENT (SBIRT/SABIRT)

- <u>Screening</u> universal screening for substance use and impact of that use
- $\begin{tabular}{ll} \hline [Assessment use of validated assessment tool to determine diagnosis and severity] \\ \hline \end{tabular}$
 - Alcohol Use Disorders Identification Test (AUDIT)

 - Drug Abuse Screening Test (DAST-10)
 Alcohol, Smoking and Substance Involvement Screening Test (NM-ASSIST)
- <u>Brief intervention</u> use of motivational interviewing concepts to reduce problematic substance use
- Referral to treatment referral to specialty substance use treatment or, in some cases, simply referral to continued assessment and follow up with their primary provider



HEALTH MANAGEMENT ASSOCIATES

S(A)BIRT

S(A)BIRT FLOW



HEALTH MANAGEMENT ASSOCIATES

111

112

CHATTER FALL

We will have two Chatter fall questions.

Please take a minute to type your response in the Zoom Group Chat, but don't click enter.

What have been the biggest challenges to implementation of screening in your setting?

What strategies have you used to overcome those challenges?

Type your response and don't click enter.

HEALTH MANAGEMENT ASSOCIATES

Stare from Unsplash



Assessment

OBJECTIVES OF A BIO-PSYCHO-SOCIAL ASSESSMENT (BPS)

- + A comprehensive biopsychosocial assessment provides:
 - + Insight into the patient's past and current life experience
 - + Provides data to make an accurate (preliminary) diagnosis
 - + An opportunity to build rapport with the patient
 - + Provides information needed to make an accurate level of care determination
 - + American Society of Addiction Medicine (ASAM) Level of Care criteria cover biopsychosocial

HEALTH MANAGEMENT ASSOCIATES

Copyright © 2021 Health Management Associates, Inc. All rights reserved.

Picture for

113

114

Assessment

BIOPSYCHOSOCIAL ASSESSMENT

- + A comprehensive biopsychosocial assessment includes:
 - + General information (housing status including who live with, religious affiliation, referral source, insurance)
 - + Medical information (past/present medical conditions, medications, surgeries, childbirths, hospitalizations)
 - + Education and Employment (highest grade, difficulty in school, past and current employment, income (legal and illegal), dependents, Social Security Benefits/Disability Benefits (SSI/SSDI), date of last employment, skill trade or technical education)
 - Legal (past and current legal issues, arrests, charges, convictions, DUI, other driving offenses, incarceration time)

HEALTH MANAGEMENT ASSOCIATES

Assessment

BIOPSYCHOSOCIAL ASSESSMENT CONT.

- Psychological (Mini mental status exam; current and past medications, inpatient and outpatient treatment, anxiety, depression, hallucinations, suicidal or homicidal)
- Family and Social (who raised, siblings, past and current relationship with family, family with past/current SUD and Department of Corrections (DOC), children, partner (with SUD?), friends and supports, hobbies, spirituality, marital status
- + SUD (substance(s)) first used and date of first use, how many days used in past 30, lifetime use and route of administration of every substance
 - + Substance(s) of choice, date of last use, overdose, and/or delirium tremens (DTs)
 - + If yes, how many times
 - + Assess use of safe drug practices
 - SUD treatment type and level of care (past, current, and # of times), MAT (past, current and # of times, if currently on buprenorphine or methadone), \$\$ spent on substances in last 30 days
- + Examples of evidence-based assessment tools:
 - NIDA Modified Assist –(not a BPS) –public domain
 Brief Addiction Monitor (BAM) public domain
 - Brief Addiction Monitor (BAM) public domain
 Addiction Severity Index public domain
- + Addiction Severity Index public dor HEALTH MANAGEMENT ASSOCIATES

116

Assessment, Level of Care

THE AMERICAN SOCIETY OF ADDICTION MEDICINE (ASAM) CRITERIA: MAPPING ASSESSMENT TO PLACEMENT

A thorough BPS assessment provides this information

Level of Care (LOC) Determination

ABBREVIATED EXAMPLE OF ASAM ADULT LEVELS OF CARE

Levels of Care Dimension	1. OUTPT	2. INTENSIVE OUTPT	3. MED MON INPT	4. MED MGD INPT
Acute Intoxication and/or Withdrawal Potential	No Risk	Minimal	Some risk	Severe risk 24- hr acute
Biomedical Conditions and Complications	No Risk	Manageable	Medical monitoring required	Medical Care Required
Emotional, Behavioral, or Cognitive Conditions and Complications	No Risk	Mild severity	Moderate	24-hour psych & addiction To required
Readiness to Change	Cooperative	Cooperative but requires structure	High resistance, needs 24-hour motivating	
Relapse, Continued Use, or Continued Problem Potential	Maintains abstinence or controls use	More symptoms, needs close monitoring	Unable to control use in outpatient care	
Recovery/Living Environment	Supportive	Less support with structure can cope	Danger to recovery. Logistical incapacity for outpatient	

HEALTH MANAGEMENT ASSOCIATES

117

HEALTH MANAGEMENT ASSOCIATES

118

Level of Care (LOC) Determination

+ ASAM Criteria is Gold Standard

- + CONTINUUM® and Co-triage® tool
- + Criteria are required in assessment tools used by providers
- + Complete for high/severe assessments
- + Available online
- + Done by RN, LCSW, PA/NP, or MD/DO
- + Part of SBIRT payment

HEALTH MANAGEMENT ASSOCIATES

ASAM and Level of Care (LOC) Determination in MN

MN Medicaid Section 1115 Waiver and related legislation requires that providers of SUD services use the ASAM Level of Care criteria

- + The legislation codifies required service standards for participating providers that are consistent with ASAM criteria
- + "All 87 Minnesota counties, 11 American Indian Tribes, and eight managed care organizations (MCOs) are required to conduct an assessment that incorporates the six dimensions of the ASAM placement criteria"
 - + Risk rating
 - + Narrative summary supporting the rating
 - + Determination of SUD diagnosis
 - + Info relevant to treatment service planning

HEALTH MANAGEMENT ASSOCIATES

119 120

hin the five broad levels of care (0.5, 1, 2, 3, 4), decimal nu

REFLECTING A CONTINUUM OF CARE

DEPARTMENT OF HUMAN SERVICES

ASAM and Level of Care (LOC) Determination in MN (cont.)

- + Providers enrolled in the 1115 demonstration evaluating use of the criteria must be compliant with the ASAM-based Standards by June 30, 2021
- + Other requirements
 - + Comprehensive assessment (c/w with ASAM criteria)
 - + Assessment summary within 3 calendar days after service initiation (or same day if comprehensive assessment is used to authorize services)
 - + Initial Services Plan
- + The state is planning to implement a requirement that residential SUD treatment facilities offer MAT services onsite or facilitate MAT access offsite at an appropriate facility.

HEALTH MANAGEMENT ASSOCIATES

121

122

ASAM Criteria Level of Care - Other Tree

ASAM ASAM Dimension 1 nos

Level of Care (LOC) Transitions

CONTINUED SERVICE CRITERIA (ASAM CRITERIA)

Retain at the present level of care if:

Making progress, but not yet achieved goals articulated in individualized treatment plan. Continued treatment at present level of care necessary to permit patient to continue to work toward his or her treatment goal

Not yet making progress but has capacity to resolve his or her problems. Actively working on goals articulated in individualized treatment plan. Continued treatment at present level of care necessary to permit patient to continue to work toward his or her treatment goals; and/or

New problems identified that appropriately treated at present level of care. This level is least intensive at which patient's new problems can be addressed effectively.

HEALTH MANAGEMENT ASSOCIATES

Level of Care (LOC) Transitions (cont.)

TRANSFER/DISCHARGE SERVICE CRITERIA (ASAM CRITERIA)

Transfer or discharge from present level of care if s/he meets the following criteria:

- Has achieved goals articulated in his or her individualized treatment plan, thus resolving problem(s) that justified admission to current level of care or
- Has been unable to resolve problem(s) that justified admission to present level of care, despite amendments to treatment plan. Treatment at another level of care or type of service therefore is indicated.
- Has demonstrated lack of capacity to resolve his or her problem(s).

 Treatment goals might be better achieved at another level of care or type of service.

124

Has experienced intensification of his or her problem(s), or has developed new problem(s), and can be treated effectively only at a more intensive level of care.

Source: The ASAM Criteria, 2013, p.303 HEALTH MANAGEMENT ASSOCIATES

31

Level of Care (LOC) Determination: Summary

- · Comprehensive assessment requires evaluation of all 6 Dimensions
- Additional assessments better positions us to fully address a client's needs
- Treatment planning can begin before LOC determination
- The LOC might have to change based on availability, but it doesn't mean we can't get started
- Where someone gets care really matters
- Implementation of ASAM assessment criteria is evolving in MN

HEALTH MANAGEMENT ASSOCIATES

125

References

REFERENCES FOR SCREENING, ASSESSMENT AND LOC DETERMINATION

- · ASAM, (2020) National Practice Guidelines for the Treatment of OUD
- MINNESOTA Statutes Minnesota Statutes, Chapter 245A. 09, Subdivision 7, paragraph (e); chapter 245G
- Center for Adolescent Substance Abuse Research, Children's Hospital of Boston. The CRAFFT screening interview. Boston (MA) CeASAR; 2009.
- Chasnoff, I. J., McGourty, R. F., Bailey, G. W., Hutchins, E., Lightfoot, S. O., Pawson, L. L., ... & Campbell, J. (2005). The 4P's Plus® screen for substance use in pregnancy: clinical application and outcomes. Journal of perinatology, 25(6), 368.
- JA Ewing "Detecting Alcoholism: The CAGE Questionnaire" JAMA 252: 1905-1907, 1984
- Reisfieled, G. et al., The ASAM Essentials of Addiction Medicine (2020), Practical Considerations in Drug Testing; p 659-66
- Laskar, D., Chemistry Toxicology General, Editor Zynger, D., 4-2019
- The Clinical Toxicology Laboratory, AACC Press, 2003, pp. 491-2
- · Broussard L, Handbook of Drug Monitoring Methods, Humana Press, 2007
- Smith, H. Opioid Metabolism (July, 2009). Mayo Clinic Proceedings; 84(7): 613–624. www.ncbi.nlm.nih.gov/pmc/articles/PMC2704133

 HEALTH MANAGEMENT ASSOCIATES

 Coppute 2211 and Management and Management Associates.

126

Group Discussion

What concepts from today's discussion about the science of SUD, screening, assessment and LOC determination will stick with you (any "ah ha" moments?) and how can you put that to use in your work?

Use the "raise your hand" feature in Zoom or simply come off mute.



HEALTH MANAGEMENT ASSOCIATES

127



Next Steps

- Join us for Session 2 next Wednesday!
- Your registration should have included a reoccurring calendar invite for all four sessions
- Please complete the evaluation for this session that will be sent out after via email (those requests CEU/CME must complete the evaluations).

Follow-up questions? cmcintire@healthmanagement.com

HEALTH MANAGEMENT ASSOCIATES

opyright © 2021 Health Management Associates, Inc. All rights reserved.