

# HMA

## The Intersection of HIV and Substance Use: Enhancing the Care Continuum with Evidence-Based Practices

Training Series: Session 3  
November 15, 2023


Copyright © 2023 Health Management Associates, Inc. All rights reserved. The content of this presentation is PROPRIETARY and CONFIDENTIAL to Health Management Associates, Inc. and only for the information of the intended recipient. Do not use, publish or redistribute without written permission from Health Management Associates, Inc.

1


## UTILIZING ZOOM

- » Ensure your audio is linked to your Zoom participant ID.
  - » If you joined the audio by computer microphone and speaker, then you're all set.
  - » If you joined the audio with a phone and did not enter your unique participant ID then enter **#<your participant ID>#** on your phone now.
 


*Note: Your unique participant ID can be found by clicking on the lower left corner of your Zoom screen where it says, 'Join Audio' and your Participant ID will appear.*
- » Ensure you are on MUTE and your camera is ON. On the bottom left corner of your screen, you will see a red line through the microphone
 




ON MUTE




Microphone ON



Camera OFF






Camera ON
- » Your participation throughout today via chat is appreciated!
  - » Locate the chat box. On the bottom middle of your screen, click on the chat icon. This will open the "Zoom Group Chat" pane on the right side of your screen. You will see messages throughout the webinar on there. When prompted by the presenters, type in your answers or questions there.



© 2023 Health Management Associates, Inc. All Rights Reserved.
HEALTH MANAGEMENT ASSOCIATES
2

2

## UTILIZING ZOOM

- » If you would like to enable closed captions during this session, please follow the steps below.
  - » On the Zoom room toolbar, tap the **Captions**  icon.
    - » You may need to tap the **More**  icon first to see the option.
  - » Ensure that the **Show Captions** toggle  is enabled.
- » If you have any issues or questions about this feature, message Carlos Mena in the chat and she can assist you.



3

## HOUSEKEEPING

Today is Session 3

Please complete the evaluation and post-test for the webinar that will be sent out via email after each session.

You will be receiving a PDF of today's presentation.

This session is being recorded.

### Follow-up questions?

Contact Carlos Mena:  
[cmena@healthmanagement.com](mailto:cmena@healthmanagement.com)

4

## CEUS AND CMES ELIGIBILITY AND DISTRIBUTION

- » This series is eligible for both CEUs and CMEs
  - » These activities have been approved for CEUs by the Minnesota Board of Behavioral Health and Therapy for 3 hours of credit for LADCs and LPC/LPCCs (total of 12 hours if all four sessions are fully attended)
  - » These activities have been approved for CMEs by the American Academy of Family Physicians for 3 hours of credit (total of 12 hours if all four sessions are fully attended)
- » To qualify for CEUs or CMEs, you are required to
  1. Complete the pre-training quiz
  2. Be in attendance for the entire session
  3. Complete the accompanying evaluation survey for each session attended
  4. Complete the post-training quiz
- » CEU/CME certificates will be issued approximately 1-2 weeks AFTER the completion of the series (Session 4: November 22th).

### Follow-up questions?

Contact Carlos Mena :  
cmena@healthmanagement.com

5

## ACKNOWLEDGMENTS



We would also like to thank our community partners for their support in developing this curriculum.



6

## LAND ACKNOWLEDGMENT



Every community owes its existence and vitality to generations from around the world who have contributed their hopes, dreams, and energy to making the history that led to this moment. Some were brought here against their will, some were drawn to leave their distant homes in hope of a better life, and some have lived on this land for more generations than can be counted. Truth and acknowledgment are critical to building mutual respect and connection across all barriers of heritage and difference.

We begin this effort to acknowledge what is buried by honoring the truth. We are standing on the ancestral lands of the Dakota people. We want to acknowledge the Dakota, the Ojibwe (pronounced ow-jeeb-way), the Ho Chunk, and the other nations of people who also call this place home. We pay respects to their elders past and present.

Please take a moment to consider the treaties made by the Tribal nations that entitle non-Native people to live and work on traditional Native lands. Consider the many legacies of violence, displacement, migration, and settlement that bring us together here today. Please join us in uncovering such truths at any and all public events.\*

\*This is the acknowledgment given in the USDAC Honor Native Land Guide – edited to reflect this space by Shannon Geshick, MTAG, Executive Director Minnesota Indian Affairs Council

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

7

7

## TODAY'S PRESENTERS



**Charles Robbins, MBA**  
(he/him/his)  
Principal  
Health Management Associates



**Linda Follenweider, MS, APRN**  
(she/her/hers)  
Managing Director  
Health Management Associates



**Shannon Robinson, MD**  
(she/her/hers)  
Principal  
Health Management Associates

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

8

8

## DISCLOSURES

Faculty	Nature of Commercial Interest
Charles Robbins, MBA	Mr. Robbins discloses that he is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients.
Linda Follenweider, MS, APRN	Ms. Follenweider discloses that she is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients.
Shannon Robinson, MD	Dr. Robinson discloses that she is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients.
Jeanene Smith, MD	Dr. Smith discloses that she is an employee of Health Management Associates, a national research and consulting firm providing technical assistance to a diverse group of health care clients.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

9

9

## AGENDA FOR WEBINAR SERIES

Session	Topics
<b>#1</b> <b>WEDNESDAY, NOV 1</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> Understanding HIV <input type="checkbox"/> HIV Testing and Treatment <input type="checkbox"/> The Science of Addiction <input type="checkbox"/> Screening, and Assessment
<b>#2</b> <b>WEDNESDAY, NOV 8</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> Ethical and Legal Issues <input type="checkbox"/> Funding and Policy Considerations <input type="checkbox"/> HIV Risk Reduction <input type="checkbox"/> SUD Harm Reduction <input type="checkbox"/> HIV and Stigma <input type="checkbox"/> Motivational Interviewing
<b>#3</b> <b>WEDNESDAY, NOV 15</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> <b>Working with Justice Involved Persons</b> <input type="checkbox"/> <b>Substance Use Disorder Treatment with Medications</b> <input type="checkbox"/> <b>Mental Health Treatment and Counseling</b> <input type="checkbox"/> <b>Stimulant Use</b> <input type="checkbox"/> <b>Chem Sex</b>
<b>#4</b> <b>WEDNESDAY, NOV 22</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> Cultural, Racial and Sexual Identities <input type="checkbox"/> Pregnancy and HIV, SUD/ODU <input type="checkbox"/> Accessing, Obtaining, and Integrating Services for Individuals with HIV and SUD in Minnesota

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

10

10

## TIME FOR A POLL

**Please indicate the sector(s) in which you currently serve:**

- A. Community based organizations (Social Services, HIV, LGBT, etc.)
- B. Corrections (includes Probation, Jail, Prison)
- C. County Behavioral Health, Public Health, Human Services
- D. Non-county behavioral health
- E. Federally Qualified Health Center (FQHC)
- F. Narcotic Treatment Program/Opioid Treatment Program
- G. Outpatient Treatment Program
- H. Residential Treatment Program
- I. Aftercare services (e.g., sober living, other recovery housing, recovery community centers, etc.)
- J. Harm Reduction Services/SSPs
- K. Other (please specify in the chat)

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

11

11

## TIME FOR A POLL

**Please indicate your primary role or discipline:**

- A. Physicians, Physician Assistant, Nurse Practitioners, Nurses (RN, LVN)
- B. Social Workers
- C. Addiction Counselors (LADCs)
- D. Peer Recovery Support Positions
- E. Substance Use Navigators (SUNs)
- F. Administrators, Program Managers
- G. Psychologists, LMFTs
- H. Criminal Justice Professionals
- I. Community Members
- J. Other (please specify in the chat)

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

12

12

## GLOSSARY OF TERMS (REVISITED)

- » Sexual orientation – a person’s identity in relation to the gender or genders to which they are sexually attracted (straight, gay, lesbian, asexual, bisexual, pansexual)
- » Gender identity and/or expression - internal perception of one’s gender; how one identifies or expresses oneself.
  - » Cisgender – a term used to describe a person whose gender identity aligns with those typically associated with the sex assigned to them at birth
  - » Transgender – refers to an individual whose current gender identity and/or expression differs from the sex they were assigned at birth (may have transitioned or be transitioning in how they are living)
  - » Gender Expansive - refers to an individual who expresses identity along the gender spectrum (genderqueer, gender nonconforming, nonbinary, agender, two spirit)
- » Sexual Minority – refers to a group whose sexual identity orientation or practices differ from the majority of and are marginalized by the surrounding society.

SOURCE: Centers for Educational Justice and Community Engagement, UC Berkeley

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

13

13

## GLOSSARY OF TERMS (REVISITED)

- » Race - is usually associated with inherited physical, social and biological characteristics. In this context that means race is associated with biology. Institutionalized in a way that has profound consequences (White, African American, American Indian Alaskan Native, Native Hawaiian or Pacific Islander)”
- » Ethnicity - a term used to categorize a group of people with whom you share learned characteristics and identify according to common racial, national tribal, religious, linguistic, or cultural origin or background. (Hispanic, Non-Hispanic Black, Non-Hispanic Black, etc.)

SOURCE: US Office of Management and Budget: Federal Register Vol. 62(210): 58782

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

14

14

## COMMON ACRONYMS (REVISITED)

ART – Antiretroviral therapy

AUD – Alcohol use disorder

IDU – Injection or intravenous drug use

MSM – Men who have sex with men

OUD – Opioid use disorder

PEH – Person(s) experiencing homelessness

PEP – Post-exposure prophylaxis

PrEP – Pre-exposure prophylaxis

PLWH – Person(s) living with HIV

PWID – Person(s) who injects drugs

SUD – Substance use disorder

15

## WORKING WITH JUSTICE-INVOLVED INDIVIDUALS

16



## LEARNING OBJECTIVES: WORKING WITH JUSTICE-INVOLVED INDIVIDUALS

I

Describe the importance of substance use disorder treatment with medications in criminal justice settings

II

List 3 actions to take to ensure continuity of care for clients upon release from justice settings

III

Compare and contrast FDA approved medications for Alcohol Use Disorder (AUD), Opioid Use Disorder (OUD), and opioid reversal

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

17

17

### How many Minnesota residents are locked up and where?

17,500 of Minnesota's residents are locked up in various kinds of facilities



Facility Type	Number of Residents
State Prisons	8,000
Local Jails	5,900
Federal Prisons	1,900
Youth	950
Involuntary Commitment	870
Indian Country	50

**PRISON**  
POLICY INITIATIVE


Details may not add to total due to rounding.  
Sources and data notes: See [www.prisonpolicy.org/reports/correctionalcontrol2023.html](http://www.prisonpolicy.org/reports/correctionalcontrol2023.html)

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

18

18



19




20

Race/Ethnicity	Count	Percentage of incarcerated population (%)	<b>MINNESOTA DEPARTMENT OF CORRECTIONS ADULT PRISON POPULATION SUMMARY (AS OF 07/01/2022)</b>  <b>Average age: 39.2</b> <b>Average ADP 2022: 7,527</b> <b>Males: 7,332 (93.6%)</b> <b>Females: 501 (6.4%)</b>
White	4,010	51.2%	
Black	2,782	36.7%	
American Indian	725	9.3%	
Asian	205	2.6%	
Unknown/ Other	21	0.3%	
<b>Total</b>	<b>7,833</b>	<b>100%</b>	

**Note:** 425 (5.4%) of the above are of Hispanic ethnicity.

[https://mn.gov/doc/assets/Adult%20Prison%20Population%20Summary%207-1-2022\\_tcm1089-534656.pdf](https://mn.gov/doc/assets/Adult%20Prison%20Population%20Summary%207-1-2022_tcm1089-534656.pdf)  
© 2023 Health Management Associates, Inc. All Rights Reserved. HEALTH MANAGEMENT ASSOCIATES

21



22

## MINNESOTA DEPARTMENT OF CORRECTIONS ADULT PRISON POPULATION SUMMARY (AS OF 07/01/2023)

Top Six Offenses	Count	Percentage (%)
Homicide	1,588	19.2%
Criminal Sexual Conduct	1,5152	18.8%
Drugs	1,331	16.1%
Weapons	797	9.6%
Assault	748	9.0%
DWI	383	4.6%
<b>Note:</b> Percentages are based on the total population of 8,274		
Releases (FY2023)	Count	Percentage (%)
Supervised Release/Parole	3,650	78.4%
Community Programs	656	14.1%
Discharge	306	6.6%
Other	43	0.9%
<b>Total</b>	<b>4,655</b>	<b>100.0%</b>

[https://mn.gov/doc/assets/Adult%20Prison%20Population%20Summary%207-1-2023\\_tcm1089-589994.pdf](https://mn.gov/doc/assets/Adult%20Prison%20Population%20Summary%207-1-2023_tcm1089-589994.pdf)

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES


23

23

## UNITED STATES DRUG OVERDOSE DEATHS THROUGH MAY 2023

Based on data available for analysis on: October 1, 2023

**Figure 1a. 12 Month-ending Provisional Counts of Drug Overdose Deaths: United States**



**United States**

May 2023: 12 month 106,539

May 2022: 12 month 107,419

<https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES


24

24

## OVERDOSE DEATH MINNESOTA

Based on data available for analysis on: October 1, 2023

Figure 1a. 12 Month-ending Provisional Counts of Drug Overdose Deaths: Minnesota



**Minnesota**

May 2023: 12 month 1,404  
 May 2022: 12 month 1,344

<https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm>

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

25

25

## HIV IN PRISON

An estimated 11,940 persons in the custody of state and federal correctional authorities were known to be living with HIV, a decline of nearly 16% from yearend 2019 (14,180).



Note: Between one and four jurisdictions did not report the number of persons living with HIV in each year of the 30-year period from 1991 to 2020. Data were imputed for those jurisdictions not reporting data using various methods; therefore, numbers presented are estimates. See Methodology. See appendix table 1 for estimates.  
 Source: Bureau of Justice Statistics, National Prisoner Statistics, 1991-2020.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

[HIV in Prisons, 2020 - Statistical Tables \(ojp.gov\)](#) 26

26

PERSON INCARCERATED WITH HIV

There are far fewer HIV-positive people in prison than in the past, but the HIV rate in prison is still triple the total U.S. rate

NUMBER OF PEOPLE IN STATE AND FEDERAL PRISONS WITH HIV, 1991-2021

Year	Number of People
1991	17,551
2001	25,757
2021	11,810

CURRENT PREVALENCE RATES OF HIV IN THE U.S. AND IN U.S. PRISONS, PER 100,000 PEOPLE

Category	Rate per 100,000
U.S. total population living with HIV, 2019 (aged 13 and older)	380
U.S. prison population living with HIV, 2021	1,144

Sources: Bureau of Justice Statistics, *HIV in Prisons* series, 1991-2021 and CDC, *Monitoring selected national HIV prevention and care objectives by using HIV surveillance data— United States and 6 dependent areas, 2019*  
[https://www.prisonpolicy.org/blog/2023/06/01/hiv\\_in\\_prisons](https://www.prisonpolicy.org/blog/2023/06/01/hiv_in_prisons)  
 Bureau of Justice Statistics, *Census of Jails, 2019*; and *Annual Survey of Jails, 2020*

PRISON POLICY INITIATIVE

© 2023 Health Management Associates, Inc. All Rights Reserved. 27

27

BURDEN OF SUBSTANCE USE DISORDER (SUD) IN CARCERAL SETTINGS

- » 11% of 18-25 year olds, and 6% of those over 25 years old have a SUD\*
- » 63% of people in jail and 58% in prison have a SUD.\*
- » Historically jails withdrew people from medication for addiction treatment.\*\*
- » Outcomes are much better if people are continued on treatment.\*\*
- » 77% of deaths within 2 weeks of release are related to overdose.
- » This can be decreased by 60-80% with access to medication \*\*\*


\*<https://www.samhsa.gov/criminal-juvenile-justice/about>  
 \*\*Rich 2015; \*\*Kimlock 2009.  
 \*\*\*Green 2019; Lim 2023

GUIDELINES FOR MANAGING SUBSTANCE WITHDRAWAL IN JAILS A Tool for Local Government Officials, Jail Administrators, Correctional Officers, and Health Care Professionals. June 2023

© 2023 Health Management Associates, Inc. All Rights Reserved. HEALTH MANAGEMENT ASSOCIATES 28

28

## DECREASE IN JAIL POPULATION DOES NOT EQUAL DECREASE IN BURDEN OF DISEASE FOR CARCERAL SETTING




Note: "Ave. Daily P" refers to average daily physical health conditions and "Ave. Daily M" refers to average daily mental health conditions.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

29

29



© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

30

30

## TRANSITION OF CARE


- » Transition of Care – The movement of a patient from one setting of care (hospital, ambulatory primary care practice, ambulatory specialty care practice, long-term care, home health, rehabilitation facility) to another.
- » Important Issues Regarding Release
  - » Narcan on release
  - » Warm handoff to community provider
  - » Challenges in jails and beyond
    - » No clear discharge date/time
    - » Release not correlated to clinical condition
    - » Housing options frequently suboptimal in supporting recovery
    - » Variability in provision of substance use disorder treatment with medications

<https://store.samhsa.gov/sites/default/files/d7/priv/sma16-4998.pdf>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

Photo Source :  
Ashley Jurius on  
Unsplash



31

## COMMUNITY OPPORTUNITIES TO MINIMIZE INCARCERATION

- » Early identification of individuals with mental and substance use disorders at all points of contact with the justice system – pre-arrest, booking, adjudication, reentry.
- » Use of screening and assessment to ensure linkage with evidence-based treatment, services and supports.
- » Diversion of individuals from the justice system into home- and community-based treatment.
- » Engaging law enforcement, first responders, and crisis management teams, justice court personnel, and community treatment providers in diversion strategies that meet both clinical and public safety needs.

<https://store.samhsa.gov/sites/default/files/d7/priv/sma16-4998.pdf>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

32

32



## COMMUNITY OPPORTUNITIES TO MINIMIZE INCARCERATION CONT.

- » Provision of training and technical assistance for law enforcement officers, juvenile and family court judges, probation officers, and other judicial decision-makers on behavioral health issues; and conversely, training for behavioral health treatment providers on criminogenic risk and the criminal and juvenile justice system.
- » Provision of an array of services and supports to enable successful reentry into the community for those transitioning from incarceration or detention including housing.
- » Assurance of equitable opportunities for diversion and linkage to community services and supports for all populations in order to decrease disproportionate minority contact with the justice system.
- » Promotion of cross-sector collaboration to better serve these populations dually involved with the behavioral health and criminal justice systems.

<https://store.samhsa.gov/sites/default/files/d7/priv/sma16-4998.pdf>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

33

33

## TIME FOR A POLL

**Statement: My organization has an active working process to identify and provide a soft landing into the community for patients with complex care management needs related to addiction and HIV upon release from carceral settings.**

- A. Yes
- B. No
- C. Not Sure

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

34

34


# SUBSTANCE USE DISORDER TREATMENT WITH MEDICATIONS

35

## WHAT IS SUBSTANCE USE DISORDER TREATMENT WITH MEDICATIONS?

- » The use of FDA-approved prescription medications, usually in combination with counseling and behavioral therapies, to provide a whole-person approach to the treatment of substance use disorders (SUD).
- » When discussing medication for opioid use disorder this is frequently referred to as Medications for Opioid Use Disorder (MOUD).
- » MOUD has proven clinically effective to alleviate symptoms of withdrawal & reduce cravings. MOUD maintenance has been proven to cut overdose rates in half and decrease rates of HIV and hepatitis C transmission.
- » Research shows that a combination of MOUD and behavioral therapies is a successful method to treat OUD.

36



37

## WHY IS MOUD IMPORTANT?

Treat Withdrawal	Address Dopamine Depletion	Treat OUD/Achieve Results
<ul style="list-style-type: none"> <li>Muscle pain, dilated pupils, nausea, diarrhea, abdominal cramping, piloerection</li> <li>Lasts 3-14 days</li> <li>Methadone or buprenorphine are recommended over abrupt cessation due to risk of return to use, overdose (OD) &amp; death</li> </ul>	<ul style="list-style-type: none"> <li>Reward/motivation pathway abnormalities persists for months after people stop using</li> <li>Treated with methadone or buprenorphine</li> </ul>	<ul style="list-style-type: none"> <li>Abstinence based treatment results in 85% using opioids within 1 year</li> <li>MOUD decreases                             <ul style="list-style-type: none"> <li>Use</li> <li>Craving</li> <li>Complications from IVDU</li> <li>Criminal behavior</li> </ul> </li> <li>MOUD increases retention in treatment</li> </ul>

Sources: ASAM, (2020) National Practice Guidelines for the Treatment of OUD, Mattick, RP & Hall W (1996) Lancet 347: 8994, 97-100. Mattick, RP et al. (2008) Cochrane Systematic Review. Mattick, RP, et al. (2009) Cochrane Systematic Review. Lobmaier, P et al. (2008) Cochrane Systematic Review. Krupitsky et al. (2011) Lancet 377, 1506-13. Kakko et al. (2003) Lancet 361(9358),662-8. Rich, JD, et al. (2015) Lancet

© 2023 Health Management Associates, Inc. All Rights Reserved. HEALTH MANAGEMENT ASSOCIATES 38

38

## FDA APPROVED MEDICATION FOR OUD

**Agonist Treatment**  
 (turns on the receptor):

- Methadone- approved for cough in 1940s, for OUD 1972
- Buprenorphine (Suboxone™ & Subutex™)- approved in 1981 for pain; oral approved for OUD 2002, patch, implants & injection later

**Antagonist Treatment**  
 (blocks receptor from turning on):

- Naltrexone (Revia™)- oral approved 1984; injectable (Vivitrol™)2006 for AUD, 2010 for OUD
- Naloxone- approved 1961, autoinjector 2014, nasal spray (Narcan™) 2015
- Nalmefene™ - approved nasal spray 2023

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

39

39

## HOW DO THESE MEDICATIONS WORK?

Methadone      Buprenorphine

Naltrexone/Naloxone/Nalmefene

**Heroin**


Agonist Treatment

**Heroin**

Antagonist Treatment

© 2023 Health Management Associates, Inc. All Rights Reserved.

40



41

### METHADONE: WHAT AND FOR WHOM?

- » Mu opioid receptor agonist
  - » No “ceiling effect”
- » Reaching a therapeutic dose takes time
  - » <60 mg/d is not therapeutic
  - » Typical dose 60-120 mg/d
  - » Increased frequency and daily dose required during pregnancy
- » Several significant drug-drug interactions
- » Illegal to write prescription for methadone to treat OUD unless:
  - » Narcotic Treatment Program (NTP)
  - » Covering a gap of no more than 3 days
  - » Patient is hospitalized

- Patients with greater than a year of an OUD\*
- Patients with a more severe OUD, such as injecting opioids
- Patients who have not reached tx goals with other MOUD
- Patients who would benefit from the closest follow up

\* Legislatively addressed in Omnibus bill; effective date pending

HEALTH MANAGEMENT ASSOCIATES


42


42


## METHADONE: GENERAL FEDERAL REGULATIONS



Delivered via  
observed  
dosing

Once patient is stable  
and after 6 weeks, can  
be given take-home  
doses (varies by  
state)\* 

Highly monitored in  
a Narcotics or  
Opioid Treatment  
Program setting  
(NTP/OTP) 

Many requirements  
for treating patients:  
therapy,  
toxicology\*... 

\* OUD $\geq$ 1 y requirement for methadone removed by Omnibus Bill 12.29.22, 18 months for HHS to implement; Proposed Rule <https://public-inspection.federalregister.gov/2022-27193.pdf>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

43

43

## METHADONE: EFFICACY DATA

- » Methadone resulted in 33% fewer opioid positive toxicology tests compared to those receiving no medication\*
  - » Everyone received psychosocial treatment
- » 4.4x more likely to stay in treatment \*
- » Reduced crime \*
- » Reduced infectious disease\*
- » Reduced death\*\*

Source:  
\* Mattick 2009 Cochrane Review  
\*\* Wakeman 2020 JAMA Open Network

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES



44

## BUPRENORPHINE: WHAT AND FOR WHOM?

- » Partial mu opioid agonist with ceiling effect
  - » Available alone or in combination w/naloxone
  - » Doses >32 mg don't cause greater effect
  - » Different formulations (sub-lingual [SL] buccal pill/film, injectable)
- » Greater binding affinity than full agonists
  - » Start buprenorphine when client in moderate withdrawal (to avoid causing precipitated withdrawal)
  - » Other opioids are not as effective when buprenorphine is present
  - » Typical dose is 16-24 mg/d
  - » Increased frequency and daily dose required during pregnancy
- » Fewer drug-drug interactions than methadone

Opioid use  
disorder or  
withdrawal

Patient wants  
agonist  
treatment

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

45

45

## BUPRENORPHINE REGULATIONS

### DEA X-Waiver updates

<https://www.deadiversion.usdoj.gov/pubs/docs/index.html>

X waiver no longer required  
Use standard DEA number for  
buprenorphine prescriptions

No cap on number of people treated with  
buprenorphine

Photo Source : Bryan Lopez on Unsplash  
© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

46

46

## BUPRENORPHINE EFFICACY

- » Rate of return to opioid use for persons taking placebo was 100% vs 25% for persons taking buprenorphine
- » If taking  $\geq 16$ mg buprenorphine you are 1.82 times more likely to stay in treatment than if on placebo
- » Decreased death\*



Source:  
 NIDA Medications to Treat Opioid Use Disorder Research Report Updated December 2021  
 Mattick 2014 Cochrane Review  
 \* Wakeman 2020 JAMA Open Network

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

47

47

## NALOXONE: OD REVERSAL AGENT AS HARM REDUCTION

- » Mu opioid antagonist
- » Shorter half-life & more rapid onset of action than naltrexone
- » High affinity, competitive binding & displaces agonists
- » Intranasal or intramuscular by bystander
- » May require more than one dose
- » Opioids have longer half-life than naloxone
- » Saves lives; no evidence for increasing drug use
- » Good Samaritan law in MN
- » <1% of those in need have access
- » Available over the counter


© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

48

48





49

## NALOXONE RESOURCES

- >> <https://www.health.state.mn.us/communities/opioids/opioid-dashboard/resources.html#naloxone>
- >> University of Minnesota Naloxone Resources  
<https://www.pharmacy.umn.edu/degrees-and-programs/continuing-pharmacy-education/continuing-education-courses/naloxone>
- >> Naloxone overdose training and kits free of charge. The following community-based organizations provide Naloxone overdose training and kits free of charge:
- >> [Steve Rummler HOPE Network](#)—Call 952-943-3937 or sign up for training from the [Steve Rummler HOPE Network](#).
- >> [Rural AIDS Action Network \(RAAN\)](#)—Call 320-257-3036.
- >> [Red Door Clinic](#)—Call 612-543-5555.
- >> [Indigenous Peoples Task Force](#)—Call 612-870-1723.
- >> [Lutheran Social Services](#)—Call 800-582-5260.

50

## TIME FOR A POLL

Do you know if your organization is currently prescribing (or providing) or doing any training on naloxone?

- A. Yes
- B. No
- C. I Don't Know

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

51

51

## NALTREXONE: WHAT AND FOR WHOM?

- » Mu opioid antagonist with high, competitive binding affinity
- » Does not treat withdrawal or underlying dopamine depletion
- » Must be opioid free x 7 days before starting
- » More widespread acceptance in criminal justice and “abstinence-only” communities
- » Evidence of decreased mortality is limited

Patients with a high degree of motivation (dopamine)

Patients with a history of OUD and Alcohol Use Disorder (AUD)

Patients who did not reach treatment goals with methadone or buprenorphine

Can be useful for occasional use or after discontinuation of methadone or buprenorphine

Source: Larochelle, et al. Medication for opioid use disorder after nonfatal opioid overdose and association with mortality. A cohort study. Annals of Internal Medicine. 169:3 (2018) 137-45.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

52

52

## NALTREXONE: GENERAL REGULATIONS



No Federal regulations inhibit the use

Not all BH clinics have RN to give injections



Multiple formulations:

- Pills at 25mg and 50 mg (50-100 mg for AUD)
- Long acting injectable 380mg (28-30 days)

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

53

53

## NALTREXONE: EFFICACY DATA

- » Extended Release (XR) Naltrexone  
90% opioid abstinent toxicology tests vs. 35% placebo\*
  - » Decreased incarceration\*\*
  - » Does not decrease death\*\*\*
- » XR Naltrexone vs usual care in HIV clinic\*\*\*\*
  - » Fewer days of opioid use for those on XR Naltrexone



This Photo by Unknown Author is licensed under CC BY

Source:

\*Krupitsky 2011 Lancet

\*\*Minozzi 2011 Cochrane Review

\*\*\*Wakeman 2020 JAMA Open Network

\*\*\*\* Korthuis 2022

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

54

54

## HOW LONG TO TREAT OUD?

- » Studies of all FDA approved meds for OUD indicate a risk of return to opioid use upon discontinuation of meds
- » **Year(s) post sobriety**, if making appropriate changes to decrease likelihood of future substance use, stable in recovery and life and wants to discontinue
  - » Social Support that supports recovery
    - » Active in 12 step meetings or
    - » Active in Self-Management and Recovery Training (SMART) meetings or
    - » Active in church
    - » Not living with people who are using
  - » Able to handle interpersonal conflicts without relapsing...
  - » Avoid tapering during big life transitions such as leaving incarceration, pregnancy or delivery, moving across the country, changing jobs

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

55

55

## TO TAPER OR NOT TO TAPER?

Evidence is clear that long-term or indefinite treatment with medications for OUDs is often required for effective and sustained outcomes<sup>1</sup>

In practice, successful tapers from methadone or buprenorphine typically occur in only about 15 percent of cases<sup>2,3</sup>

According to the U.S. Surgeon General, successful tapers typically occur, if at all, when individuals have been treated with Medicated Assisted Treatment (MAT) for at least 3 years<sup>4</sup>

1. National Academies of Sciences, Engineering, and Medicine. (2019). *Medications for opioid use disorder save lives*. Washington, DC: The National Academies Press.
2. Nosyk, B., Sun, H., Evans, E., Marsh, D. C., Anglin, M. D., Hser, Y. I. et al. (2012). Defining dosing pattern characteristics of successful tapers following methadone maintenance treatment: Results from a population-based retrospective cohort study. *Addiction*, 107, 1621-1629.
3. Substance Abuse and Mental Health Services Administration. (2018). *Medications for opioid use disorder: Treatment improvement protocol (TIP 63) for healthcare and addiction professionals, policy makers, patients and families*. (Rep. No. HHS Publication No. SMA 18-5063). Bethesda, MD: Author.
4. Substance Abuse and Mental Health Services Administration and Office of the Surgeon General. (2018). *Facing addiction in America: The Surgeon General's spotlight on opioids*. Washington, DC: US Department of Health and Human Services.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

56

56

## REFERENCES: OUD MEDICATION

- » The ASAM National Practice Guideline for the Treatment of Opioid Use Disorder: 2020 Focused Update. (2020) *J Addict Med* 14(2S Suppl 1):1-91. doi: 10.1097/ADM.0000000000000633. Erratum in: *J Addict Med*. 2020 May/June;14(3):267. PMID: 32511106.
- » <https://www.deadiversion.usdoj.gov/pubs/docs/index.html> Eliminating the X Waiver
- » [www.druginserter.com](http://www.druginserter.com)
- » Green, T. C. et al. (2018). Postincarceration Fatal Overdoses After Implementing Medications for Addiction Treatment in a Statewide Correctional System. *JAMA psychiatry*, 75(4), 405–407. <https://doi.org/10.1001/jamapsychiatry.2017.4614>
- » [Healthresearchfunding.org](https://healthresearchfunding.org)(2019) <https://healthresearchfunding.org/24-opiate-addiction-recovery-statistics/> 24 Shocking Opiate Addiction Recovery Statistics
- » Korthis PT, et. al. HIV clinic-based extended-release naltrexone versus treatment as usual for people with HIV and opioid use disorder: a non-blinded, randomized non-inferiority trial. *Addiction*. 2022 Jul;117(7):1961-1971. doi: 10.1111/add.15836. Epub 2022 Mar 2. PMID: 35129242; PMCID: PMC9314106.
- » Krupitsky, et. al. Injectable extended-release naltrexone for opioid dependence: a double-blind placebo controlled, multicenter randomized trial. 2011; *Lancet* 377: 1506-13.
- » Kakko et al. 1-year retention and social function after buprenorphine-assisted relapse prevention treatment for heroin dependence in Sweden: a randomised, placebo-controlled trial. *Lancet* (2003) 361(9358):662-8.
- » Laroche, et al. Medication for opioid use disorder after nonfatal opioid overdose and association with mortality. A cohort study. *Annals of Internal Medicine*. 169:3 (2018) 137-45.
- » Lim S, Cherian T, Katyal M, Goldfeld KS, McDonald R, Wiewel E, Khan M, Krawczyk N, Braunstein S, Murphy SM, Jalali A, Jeng PJ, MacDonald R, Lee JD. Association between jail-based methadone or buprenorphine treatment for opioid use disorder and overdose mortality after release from New York City jails 2011-17. *Addiction*. 2023 Mar;118(3):459-467. doi: 10.1111/add.16071. Epub 2022 Nov 16. PMID: 36305669; PMCID: PMC9898114.
- » Mattick, RP, et al. (2009) Methadone maintenance therapy versus no opioid replacement therapy for opioid dependence. *Cochrane Systematic Review*.
- » Mattick, RP, et al. (2014) Buprenorphine maintenance versus placebo or methadone maintenance for opioid dependence. *Cochrane Systematic Review*.
- » Metzger DS et al., "Human Immunodeficiency Virus Seroconversion Among Intravenous Drug Users In- and Out-of-Treatment: An 18-Month Prospective Follow-Up." *Journal of Acquired Immune Deficiency Syndromes* 6, no. 9 (1993): 1049–56, <http://www.ncbi.nlm.nih.gov/pubmed/8340896>.
- » Minozzi S, Amato L, Vecchi S, Davoli M, Kirchmayer U, Verster A. Oral naltrexone maintenance treatment for opioid dependence. *Cochrane Database Syst Rev*. 2011;2011(4):CD001333. Published 2011 Apr 13. doi:10.1002/14651858.CD001333.pub4

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

57

57

## REFERENCES: OUD MEDICATION

- » NIDA Medications to Treat Opioid Use Disorder Research Report Updated December 2021.
- » *Principals of Drug Addiction Treatment: A Research Based Guide.* National Institute on Drug Abuse. Ed. NIDA International Program.
- » <https://public-inspection.federalregister.gov/2022-27193.pdf> Proposed Rules MOUD
- » Rich, JD, et al. Continuation versus forced withdrawal on incarceration in a combined US prison and jail: a randomised, open-label trial. *Lancet* (2015) 386 (9991): 350-359.
- » Substance Abuse Mental Health Services Agency. Medications for Opioids Use Disorder: For Healthcare and Addiction Professionals, Policymakers, Patients and Families. Rockville MD SAMHSA Treatment Improvement Protocol (TIP) Series, No. 63 (2021).
- » Schwartz RP, et al. Opioid agonist treatments and heroin overdose deaths in Baltimore, Maryland, 1995-2009. *Am J Public Health*. 2013 May;103(5):917-22. doi: 10.2105/AJPH.2012.301049. Epub 2013 Mar 14. PMID: 23488511; PMCID: PMC3670653
- » Treatment Research Institute (TRI), Ed. "Cost Utilization Outcomes of Opioid Dependence Treatment" *American Journal of Managed Care* 2011.
- » Tsui, Ji et al., "Association of Opioid Agonist Therapy With Lower Incidence of Hepatitis C Virus Infection in Young Adult Injection Drug Users," *JAMA Internal Medicine* 174, no. 12 (2014): 1974–81, <http://archinte.jamanetwork.com/article.aspx?articleid=1918926>
- » United States Department of Justice. Office of Justice Programs. Bureau of Justice Assistance. GUIDELINES FOR MANAGING SUBSTANCE WITHDRAWAL IN JAILS A Tool for Local Government Officials, Jail Administrators, Correctional Officers, and Health Care Professionals. June 2023
- » Wakeman, SE, et. al. (2020) Comparative Effectiveness of Different Treatment Pathways for Opioid Use Disorder. *JAMA Open Network*. 3 (2).
- » Initiative, P. P. (n.d.). Minnesota incarceration pie chart 2023. [www.prisonpolicy.org](http://www.prisonpolicy.org). Retrieved November 15, 2023, from [https://www.prisonpolicy.org/graphs/correctional\\_control2023/MN\\_incarceration\\_2023.html](https://www.prisonpolicy.org/graphs/correctional_control2023/MN_incarceration_2023.html)
- » Initiative, P. P. (n.d.). Minnesota profile. [www.prisonpolicy.org](http://www.prisonpolicy.org). <https://www.prisonpolicy.org/profiles/MN.html>
- » MINNESOTA DEPARTMENT OF CORRECTIONS. (n.d.). Retrieved November 15, 2023, from [https://mn.gov/doc/assets/Adult%20Prison%20Population%20Summary%207-1-2023\\_tcm1089-589994.pdf](https://mn.gov/doc/assets/Adult%20Prison%20Population%20Summary%207-1-2023_tcm1089-589994.pdf)
- » Initiative, P. P. (n.d.). New data on HIV in prisons during the COVID-19 pandemic underscore links between HIV and incarceration. [www.prisonpolicy.org](http://www.prisonpolicy.org). Retrieved November 15, 2023, from [https://www.prisonpolicy.org/blog/2023/06/01/hiv\\_in\\_prisons](https://www.prisonpolicy.org/blog/2023/06/01/hiv_in_prisons)
- » Bureau of Justice Statistics, *Census of Jails, 2019*; and *Annual Survey of Jails, 2020*

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

58

58

## WHY MEDICATIONS FOR ALCOHOL USE DISORDER IS IMPORTANT?

Increased retention in treatment

Decreased drinking

Decreased cravings

Decreased healthcare costs

Disulfiram

Naltrexone  
(oral and intramuscular)

Acamprosate

© 2023 Health Management Associates, Inc. All Rights Reserved. HEALTH MANAGEMENT ASSOCIATES 59

59

© Elsevier. Rang et al: Pharmacology 6e - www.studentconsult.com

## DISULFIRAM: MECHANISM OF ACTION

HEALTH MANAGEMENT ASSOCIATES 60

60

## DISULFIRAM FOR ALCOHOL USE DISORDER (AUD)

- » Approved decades ago; most recent data does NOT show overwhelming efficacy\*
- » Once per day dosing
- » Inhibits multiple P450 and other liver enzymes
- » Drug Interactions: benzodiazepines, phenytoin, pimozide, tricyclic antidepressants (TCAs), warfarin, sulfonyleureas, metronidazole, amoxicillin, isoniazid
- » Contraindications/precautions: alcohol use, hypersensitivity to rubber, severe coronary artery disease (CAD), cirrhosis, severe renal impairment, psychosis, depression, diabetes mellitus (DM), epilepsy
- » Extensively metabolized
- » Extensive list of side effects

Source: \* Garbutt JC, West SL, Carey TS, et al. Pharmacological treatment of alcohol dependence. J Am Med Assoc. 1999; 281(14):1318-1325.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

61

61

## NALTREXONE FOR AUD

Few side effects

Drug Interactions: opioids

» No P450 interactions

Contraindications: severe acute hepatitis

Well studied in mild and moderate cirrhosis

Safe in mild renal disease

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

62

62

## NALTREXONE EFFICACY

	Oral	Intramuscular
Reduced drinking days	Yes	Yes
Reduced heavy drinking days	Yes	Yes
Decreased opioid use	Yes	Yes
Decreased cravings	Yes	
Increased time to first drink	Yes	Yes
Treatment retention	Higher	Highest
Discontinuation of medication		Lower than oral
Decreased ED visits		Lower
Decreased hospitalizations		Lower
Decreased pharmacy cost		Lower
Decreased nonpharmacy costs		Lower

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

63

63

## ACAMPROSATE: MECHANISM

In someone with an active alcohol use disorder acamprosate decreases glutamate release and decreases GABA transmission



© 2023 Health Management Associates, Inc. All Rights Reserved.

64

64



## ACAMPROSATE FOR AUD

- » Effective
  - » Decreased quantity and frequency
  - » Increased retention in treatment and abstinence
- » Three times per day dosing
- » Drug Interactions: none
- » Contraindications: severe renal impairment
- » 333mg three times a day (TID) moderate renal impairment (creatinine clearance 30-50ml/m)
- » Few side effects
- » No metabolism

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

65

65

## TIME FOR A POLL

**Question: Do you know anyone on medication for AUD?**

- A. Yes
- B. No

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

66

66

## REFERENCES: AUD MEDICATION

- » Chick, J., Leher, P., & Landron, F. (2003). Does Acamprosate Improve Reduction of Drinking As Well As Aiding Abstinence? *Journal of Psychopharmacology*, 17(4), 397–402. doi:10.1177/0269881103174017
- » Garbutt, J. C., West, S. L., Carey, T. S., Lohr, K. N., & Crews, F. T. (1999). Pharmacological Treatment of Alcohol Dependence. *JAMA*, 281(14), 1318. doi:10.1001/jama.281.14.1318.
- » Kahan, M., Wilson, L., & Becker, L. (1995). Effectiveness of physician-based interventions with problem drinkers: a review. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*, 152(6), 851–859.
- » Mann, K., Leher, P., & Morgan, M. Y. (2004). The Efficacy of Acamprosate in the Maintenance of Abstinence in Alcohol-Dependent Individuals: Results of a Meta-Analysis. *Alcoholism: Clinical & Experimental Research*, 28(1), 51–63. doi:10.1097/01.alc.0000108656.81563.05
- » Solberg, L. I., Maciosek, M. V., & Edwards, N. M. (2008). Primary Care Intervention to Reduce Alcohol Misuse. *American Journal of Preventive Medicine*, 34(2), 143–152.e3. doi:10.1016/j.amepre.2007.09.035
- » Substance Abuse and Mental Health Services Administration and National Institute on Alcohol Abuse and Alcoholism, Medication for the Treatment of Alcohol Use Disorder: A Brief Guide. HHS Publication No. (SMA) 15-4907. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2015.
- » Srisurapanont, M., & Jarusuraisin, N. (2005). Naltrexone for the treatment of alcoholism: a meta-analysis of randomized controlled trials. *The International Journal of Neuropsychopharmacology*, 8(2), 267–280. doi:10.1017/s1461145704004997
- » Tidey et al. (2008) Moderators of naltrexone's effect on drinking, urge and alcohol effects in non-treatment-seeking heavy drinkers in natural environment. *Alcohol Clin Exp Res*; 32 (1): 58-66.
- » Wong, J., Saver, B., Scanlan, J. M., Gianutsos, L. P., Bhakta, Y., Walsh, J., ... & Rudolf, V. (2020). The ASAM clinical practice guideline on alcohol withdrawal management. *Journal of Addiction Medicine*, 14(3S), 1-72.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

67

67

## 5-MINUTE STRETCH BREAK!



68

# COUNSELING FOR CO-OCCURRING HIV & SUD

69

69

## LEARNING OBJECTIVES: COUNSELING FOR CO-OCCURRING HIV & SUD

- I  
Discuss coping with a HIV diagnosis and preparing patients for disclosure
- II  
Identify at least 3 considerations for mental health treatment of individuals with HIV and SUD
- III  
Distinguish acute and chronic risk of suicidality in individuals with HIV and SUD

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES


70

70

## WHY IS IT IMPORTANT TO ADDRESS SUD IN PERSONS WITH HIV?

### Substance use accelerates the progression of HIV

- » Increases viral load
- » Increases likelihood of AIDs related morbidity (even when adherent to antiretroviral medications)
- » Decreases medication adherence



Sources: Dash, 2015; Schaffer 2017; Strazza 2011; Dahal 2015; Andriote 2012; NIDA 2021 <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders/>  
Photo Source: National Cancer Institute on Unsplash

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

71

71

## WHY IS IT IMPORTANT TO ADDRESS SUD IN PERSONS WITH HIV?

### “Substances of abuse” weaken the blood brain barrier

- » Allowing HIV to more easily enter the brain
- » Allows infection and damage to nerves and supporting cells (glia)
- » Triggers release of neurotoxins
- » Can lead to dementia
  - » 50% of people with HIV have neurocognitive disorders



Sources: Dash, 2015; Schaffer 2017; Strazza 2011; Dahal 2015; Andriote 2012; NIDA 2021 <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders/>  
Photo Source: Milad Fakurian on Unsplash

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

72

72

## HIV TESTING

- » 19% of 15-44yo in the United States were tested for HIV in the past year
- » Only one-third of SUD programs offer onsite HIV testing



Sources: NIDA 2021 <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders>  
Substance Abuse and Mental Health Services Administration. (2021). Treating Substance Use Disorders Among People with HIV. Advisory  
Photo Source: Testaltize.me on Unsplash

© 2023 Health Management Associates, Inc. All Rights Reserved. HEALTH MANAGEMENT ASSOCIATES

73

## HIV TESTING RECOMMENDATIONS

- ✓ SAMHSA recommends universal HIV testing for
  - » Persons 15-65yo (and all pregnant persons)
  - » Younger and older persons at increased risk, such as:
    - » People who inject drugs
    - » People who have condomless sex
    - » People who participate in commercial sex work
- ✓ US Preventative Task Force Rating A
  - » Requires Medicare and Medicaid to pay for testing
  - » Rapid tests are available- results within 30 minutes
  - » Provide pre and post test counseling- reviewed in other talks

Sources: NIDA 2021 <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders>  
Substance Abuse and Mental Health Services Administration. (2021). Treating Substance Use Disorders Among People with HIV. Advisory.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

74

74

## STTR MODEL OF CARE

- » Testing persons who inject drugs every 6 months is cost effective
- » **Recommendation:** Inpatient and outpatient mental health settings should offer routine opt out testing to improve case finding




Chart review compared to blood samples from 2 inpatient psychiatric units:  
21% of patients with HIV positive blood samples did not have documentation of infection in medical record

Sources: NIDA 2021 <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders>  
Hutchinson AB, Farnham PG, Sansom SL, Yaylali E, Mermis JH. Cost-Effectiveness of Frequent HIV Testing of High-Risk Populations in the United States. *J Acquir Immune Defic Syndr* 1999; 20(6):71(3):323-330. doi:10.1097/QAI.000000000000038.  
Gallely CL, Pinkerton SD, Petroll AE. CDC recommendations for opt-out testing and reactions to unanticipated HIV diagnoses. *AIDS Patient Care STDS*. 2008 Mar;22(3):189-93. doi: 10.1089/apc.2007.0104. PMID: 18290754; PMCID: PMC2728134.  
Branson BM, Handsfield HH, Lampe MA, Janssen RS, Taylor AW, Lyss SB, Clark JE; Centers for Disease Control and Prevention (CDC). Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep*. 2006 Sep 22;55(RR-14):1-17. quiz CE1-4. PMID: 16988943.  
Rothbard AB, Blank MB, Slaab JP, et al. Previously Undetected Metabolic Syndromes and Infectious Diseases Among Psychiatric Inpatients. *Psychiatric services*. 2009;60(4):534-537.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

75

75

## EPIDEMIOLOGY- HIV & MENTAL HEALTH

- » Up to 70% of people living with HIV have a history of trauma
- » 54% of people living with HIV have post-traumatic stress disorder (PTSD)
- » People living with HIV are twice as likely to develop depressive symptoms compared to those at risk but who are not living with HIV
- » People living with HIV experience higher rates of depression than the general population
- » Key feature of depression, as compared to adjustment disorder or side effects from medication, is loss of pleasure

Sources: Kessler, R.C. 2005, Andriote, JM. 2012, Gaynes, B.N. 2008, Blank M.B. 2013

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES



76

## EPIDEMIOLOGY- HIV & MENTAL ILLNESS

- » Twenty-two percent (22%) of people with HIV have depression
  - » Of those 78% **ALSO** have an anxiety disorder
  - » Of those 61% **ALSO** have an SUD
- » Six percent (6%) of people with HIV have schizophrenia, as compared to 1% of the general population
- » Those with schizophrenia are **1.5x** as likely to contract HIV
- » Those with affective disorders were **3.8x** as likely to contract HIV



Sources: Kessler, R.C. 2005, Andriote, JM. 2012, Gaynes, B.N. 2008, Blank M.B.2013

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

77

77

## SUD, HIV AND MENTAL ILLNESS

- » Only 35% of people in 10 outpatient HIV clinics reported talking to primary care provider (PCP) about alcohol use
- » < 50% of providers in hospital-based HIV care programs conducted recommended screening and brief interventions for reducing alcohol
- » Substance use may increase high-risk sexual practices



Sources: Staruss, S.M. 2009  
Andriote, JM. 2012

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

78

78

## COUNSELING: COPING WITH AN HIV DIAGNOSIS

- » Coping with the diagnosis of HIV
  - » is a form of grieving
  - » is different from having a major depressive episode
  - » may require treatment
    - » support or psychotherapy
    - » will not respond to antidepressants



Sources: Andriote, JM. 2012 <http://www.aidsmap.com/news/aug-2021/hardest-outcome-all-hiv-and-suicide>  
 Photo Source :LinkedIn Sales Solutions on Unsplash

© 2023 Health Management Associates, Inc. All Rights Reserved

HEALTH MANAGEMENT ASSOCIATES

79

79

## COUNSELING RECOMMENDATIONS

1. Don't try to solve or fix things, but....
  - Housing is important
  - Social support is important
  - Medical care is important
  - These things helps establish a sense of control over one's life
2. Don't minimize someone's feelings
3. Don't tell people to pull themselves together
4. Listen... for risks and for talk of the future

Sources: Andriote, JM. 2012 <http://www.aidsmap.com/news/aug-2021/hardest-outcome-all-hiv-and-suicide>

© 2023 Health Management Associates, Inc. All Rights Reserved

HEALTH MANAGEMENT ASSOCIATES

80

80



## CONSIDERATIONS FOR MENTAL HEALTH TREATMENT OF INDIVIDUALS WITH HIV AND SUD

- » Major Depression, among those living with HIV, responds to the same treatments:
  - » Evidence-based psychotherapy
  - » Evidence-based medications
- » As with other conditions, keep drug-drug interactions in mind
- » Depression & bipolar disorder can make medication adherence challenging

**ANTIDEPRESSANT TREATMENT OF DEPRESSION  
RESULTS IN LOWER HEALTHCARE COSTS**

- » Persons with bipolar disorder and HIV are more likely to have unprotected intercourse with HIV negative partners
- » The risk of suicide is higher for those with HIV (at all stages) as compared to the general population

Sources: Andriote, JM. 2012 & Blank MB 2013

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

81

81

## SUD TREATMENT FOR THOSE LIVING WITH HIV

- » Cognitive Behavioral Therapy (CBT) & Motivational Interviewing (MI)
  - » Reduce drug use
  - » Reduce high risk sexual behaviors
  - » Reduce viral load
  - » Improve adherence to antiretrovirals
- » Medication for opioid use disorder
  - » Methadone and buprenorphine are associated with a 54% reduction in risk of HIV infection in persons who inject drugs

Source: NIDA 2021 <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders>

**SUD Treatment is HIV Prevention!**

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

82

82

## EPIDEMIOLOGY- SUICIDALITY & HIV

### Suicide

- » 3rd most common cause of death in 15-29yo women
- » 4th most common cause of death in 15-29yo men
- » No relationship to income
- » A life-threatening illness is a one of the most strongly predictive factors for completed suicide
- » Suicide rate in the first year after an HIV diagnosis is 5x the rate in the general population. Suicide in the first year after an HIV diagnosis accounts for 40% of all suicide in persons with HIV.

Sources: <http://www.aidsmap.com/news/aug-2021/hardest-outcome-all-hiv-and-suicide> <https://www.health.state.mn.us/people/syringe/suicide.pdf>

Suicide Attempt Rate  
*People living with HIV: 16%*  
*General Population: 3%*

Suicidal Ideation Rate  
*People living with HIV: 23%*  
*General Population: 9%*

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

83

83

## TIME FOR A POLL

**People who talk about suicide, do not attempt suicide.**

- A. True
- B. False

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

84

84

## RISK FACTORS FOR SUICIDE



DEPARTMENT  
OF HEALTH

### Suicidal Ideation Risk Assessment

STEPS AND RESOURCES FOR EXPLORING THOUGHTS OF SUICIDE

- Trauma
- Triggering event- stressor
- Ideation & past behavior
- Health-medical, mental and substance
- Purposeless, hopeless
- Poor sleep
- Mood, anxiety, anger, withdrawal
- Reckless, impulsive

Sources: <https://www.health.state.mn.us/people/syringe/suicide.pdf>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

85

85

## ASSESSMENT FOR SUICIDALITY

- » Which factors can be modified to reduce risk?
  - » Opportunities for healing
  - » Reduce harms
- » Protective factors
  - » Connectedness
  - » Support
  - » Skills- problem solving, coping, healing


Sources: <https://www.health.state.mn.us/people/syringe/suicide.pdf>  
Photo Source: Glenn Carstens-Peters on Unsplash

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

86

86



## ASSESSMENT RECOMMENDATIONS

1. Be mindful that protective factors are unique to each person
2. Use the person's language
3. Ask open ended questions such as:
  - » What are things that keep you safe?
  - » When this occurred in the past what has stopped you?
  - » Who are the people who lift your spirits?
  - » What activities lift your spirits?
  - » What would you like to develop within yourself in the future?
4. Try to identify protective factors that can be enhanced

Sources: <https://www.health.state.mn.us/people/syringe/suicide.pdf>

87

## INTEGRATED PRIMARY HIV & BEHAVIORAL HEALTH CARE

### Benefits of Integration

- » Increases likelihood of follow through on referrals
- » Improve physical health outcomes
- » Increased savings in healthcare cost
- » Reduce emergency room use

### Ryan White HIV/ AIDS Treatment Extension Act 2009

- » Aligns with HHS guidelines
- » Mandates include:
  1. Universal depression and SUD screening
    - » MH screening rates currently are between 80%-100%
    - » SUD screening rates currently are much lower
  2. Establishment of follow up plan

88

## REFERENCES:

- » Dash S, Balasubramaniam M, Villalta F, Dash C, Pandhare J. Impact of cocaine abuse on HIV pathogenesis. *Front Microbiol.* 2015 Oct 20;6:1111. doi: 10.3389/fmicb.2015.01111. PMID: 26539167
- » Schaefer CP, Tome ME, Davis TP. The opioid epidemic: a central role for the blood brain barrier in opioid analgesia and abuse. *Fluids Barriers CNS.* 2017 Nov 29;14(1):32. doi: 10.1186/s12987-017-0080-3. PMID: 29183383
- » Strazza M, Pirrone V, Wigdahl B, Nonnemacher MR. Breaking down the barrier: the effects of HIV-1 on the blood-brain barrier. *Brain Res.* 2011 Jul 5;1399:96-115. doi: 10.1016/j.brainres.2011.05.015. Epub 2011 May 14. PMID: 21641584
- » Dahal S, Chitti SV, Nair MP, Saxena SK. Interactive effects of cocaine on HIV infection: implication in HIV-associated neurocognitive disorder and neuroAIDS. *Front Microbiol.* 2015 Sep 8;6:931. doi: 10.3389/fmicb.2015.00931. PMID: 26441868
- » Andriote. HIV and Clinical Depression APA Fact Sheet 2012, Arlington VA
- » NIDA 2021 <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders>
- » Substance Abuse and Mental Health Services Administration. (2021). *Treating Substance Use Disorders Among People with HIV. Advisory.*
- » Blank MB, Himelhoch S, Walkup J, Eisenberg MM. Treatment considerations for HIV-infected individuals with severe mental illness. *Curr HIV/AIDS Rep.* 2013 Dec;10(4):371-9. doi: 10.1007/s11904-013-0179-3. PMID: 24158425
- » Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005 Jun;62(6):593-602. doi: 10.1001/archpsyc.62.6.593. Erratum in: *Arch Gen Psychiatry.* 2005 Jul;62(7):768. PMID: 15939837.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

89

89

## REFERENCES:

- » Andriote, J. APA Fact Sheet HIV mental health treatment issues HIV and clinical depression 2012
- » Gaynes BN, Pence BW, Eron JJ Jr, Miller WC. Prevalence and comorbidity of psychiatric diagnoses based on reference standard in an HIV+ patient population. *Psychosom Med.* 2008 May;70(4):505-11. doi: 10.1097/PSY.0b013e31816aa0cc. Epub 2008 Mar 31. PMID: 18378865
- » Strauss SM, Rindskopf DM. Screening patients in busy hospital-based HIV care centers for hazardous and harmful drinking patterns: the identification of an optimal screening tool. *J Int Assoc Physicians AIDS Care (Chic).* 2009 Nov-Dec;8(6):347-53. doi: 10.1177/1545109709350509. Epub 2009 Oct 22. PMID: 19850861
- » Substance Abuse and Mental Health Services Administration and Health Resources and Services Administration, *The Case for Behavioral Health Screening in HIV Care Settings.* HHS Publication No. SMA-16-4999. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2016.
- » NIDA, 2021. <https://www.drugabuse.gov/publications/research-reports/common-comorbidities-substance-use-disorders>
- » Hutchinson AB, Farnham PG, Sansom SL, Yaylali E, Mermin JH. Cost-Effectiveness of Frequent HIV Testing of High-Risk Populations in the United States. *J Acquir Immune Defic Syndr.* 1999. 2016;71(3):323-330. doi:10.1097/QAI.0000000000000838.
- » Galletly CL, Pinkerton SD, Petroll AE. CDC recommendations for opt-out testing and reactions to unanticipated HIV diagnoses. *AIDS Patient Care STDS.* 2008 Mar;22(3):189-93. doi: 10.1089/apc.2007.0104. PMID: 18290754; PMCID: PMC2728134.
- » Branson BM, Handsfield HH, Lampe MA, Janssen RS, Taylor AW, Lyss SB, Clark JE; Centers for Disease Control and Prevention (CDC). Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep.* 2006 Sep 22;55(RR-14):1-17; quiz CE1-4. PMID: 16988643.
- » Rothbard AB, Blank MB, Staab JP, et al. Previously Undetected Metabolic Syndromes and Infectious Diseases Among Psychiatric Inpatients. *Psychiatric services.* 2009;60(4):534-537.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

90

90

# STIMULANT USE

91

91

## LEARNING OBJECTIVES: STIMULANT USE AND PERSONS WHO ENGAGE IN CHEMSEX



© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

92

92

# WHAT ARE STIMULANTS?

## » Cocaine

### » “Psychostimulants with abuse potential”

- » Mahuang, ephedra & khat- plants
- » Pseudoephedrine, ephedrine & cathinone & cathine-chemical in above plants
- » “Bath salts” (synthetic man made cathinones)
- » Amphetamine (synthetic)
  - » Methamphetamine (dextro & levo)
  - » Amphetamine (dextro & levo)
  - » MDMA/ecstasy = Molly = methylenedioxy-methamphetamine
  - » Methylphenidate = Ritalin™
- » Methylxanthines (naturally occurring)
  - » Caffeine (coffee)
  - » Theophylline (tea)
  - » Theobromine (chocolate)




© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

93

93

# DRUG SEIZURES IN MINNESOTA




Source: Minnesota Department of Public Safety, Violent Crime Enforcement Teams (VCET) Dashboard  
<https://dps.mn.gov/divisions/ojp/statistical-analysis-center/Pages/vcet-dashboards.aspx>

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

94

94



95




96



# AMPHETAMINE USE NATIONALLY & LOCALLY

**Methamphetamine Use in the Past Year among Individuals Aged 12 or Older, by Geographic Area**



Source: <https://pdas.samhsa.gov/saes/state>  
 NOTE: Estimates from 2021 are not comparable to estimates from previous years due to changes in NSDUH survey methodology.

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

97

97

# STIMULANT (COCAINE AND METHAMPHETAMINE) USE MINNESOTA

**Prevalence among Individuals Aged 12 or Older in Minnesota, by Outcome**



Source: <https://pdas.samhsa.gov/saes/state>  
 NOTE: Estimates from 2021 are not comparable to estimates from previous years due to changes in NSDUH survey methodology.



© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

98

98

## METHAMPHETAMINE EMERGENCY VISITS & HOSPITAL UTILIZATION IN THE U.S.




Source: <https://www.nihem.org/categories/beyond-opioids-rapid-increase-in-drug-deaths-involving-stimulants>

99

## DRUG RELATED EMERGENCY DEPARTMENT VISITS 2021

Figure 4.1 Estimated percentages and counts of drug-related ED visits by the top five drugs (January 2021-December 2021)

Substance Abuse and Mental Health Services Administration. (2022). Preliminary Findings from Drug-Related Emergency Department Visits, 2021; Drug Abuse Warning Network (HHS Publication No. PEP22-07-03-001). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Retrieved from <https://www.samhsa.gov/data/>




The top five drugs in all drug-related ED visits in 2021 were alcohol, opioids, methamphetamine, marijuana, and cocaine. Alcohol accounted for the majority of drug-related ED visits at 39.33 percent (2.9 million), followed by opioids at 14.07 percent (1.03 million). Opioids were further broken down into three categories—fentanyl, heroin, and other opioids. Among these opioid categories, heroin accounted for the highest percentage.

100

## STIMULANT OVERDOSE DEATHS CONTINUE TO RISE NATIONALLY AND LOCALLY

Based on data available for analysis on: October 1, 2023

After opening the drug class dropdown, click the top of the dropdown menu again to make the checkboxes disappear.




Legend for Drug or Drug Class  
 Cocaine (T43.5)  
 Psychostimulants with abuse potential (T43.6)

— Reported Value  
 ○ Predicted Value

United States

Based on data available for analysis on: October 1, 2023

After opening the drug class dropdown, click the top of the dropdown menu again to make the checkboxes disappear.



Legend for Drug or Drug Class  
 Cocaine (T43.5)  
 Psychostimulants with abuse potential (T43.6)

— Reported Value  
 ○ Predicted Value

Minnesota

Source:  
<https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm#dashboard>


HEALTH MANAGEMENT ASSOCIATES

101

101

## METHAMPHETAMINES OVERDOSES WITH AND WITHOUT OPIOIDS

Figure 7. National Overdose Deaths Involving Psychostimulants with Abuse Potential (Primarily Methamphetamine)\*, by Opioid Involvement, Number Among All Ages, 1999-2021




\*Among deaths with drug overdose as the underlying cause, the psychostimulants with abuse potential (primarily methamphetamine) category was determined by the T43.6 ICD-10 multiple cause-of-death code. Abbreviated to *psychostimulants* in the bar chart above. Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2021 on CDC WONDER Online Database, released 1/2023.

102

# COCAINE OVERDOSES WITH AND WITHOUT OPIOIDS

Figure 8. National Drug Overdose Deaths Involving Cocaine\*, by Opioid Involvement, Number Among All Ages, 1999-2021



\*Among deaths with drug overdose as the underlying cause, the cocaine category was determined by the T40.5 ICD-10 multiple cause-of-death code. Source: Centers for Disease Control and Prevention, National Center for Health Statistics, Multiple Cause of Death 1999-2021 on CDC WONDER Online Database, released 1/2023.

Source: <https://nida.nih.gov/research-topics/trends-statistics/overdose-deaths-rates/#~:text=Drug%20overdose%20deaths%20involving%20psychostimulants,to%2032%2C537%20deaths%20in%202021.>  
 © 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

103

103

## IN THE CHAT BOX PLEASE ANSWER THIS QUESTION:

*Do you prefer:*

**Coffee**

**Tea**

**Chocolate**

**Soda**

*I refuse to pick just one*



© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

104

104

## MEDICINAL USES FOR STIMULANTS

- » Cocaine- used as a vasoconstrictor & numbing agent
- » “Psychostimulants with abuse potential”
  - » Ephedra- made into pseudoephedrine and used for
    - » allergies and colds
    - » Khat used for depression, obesity, fatigue in middle east
    - » Amphetamines are used for obesity, narcolepsy & Attention Deficit Hyperactivity Disorder
  - » Methylxanthines
    - » Caffeine (coffee)
    - » Theophylline (tea) used for asthma
    - » Theobromine (chocolate)

### Amphetamine dosing:

ADHD 2.5 mg/day to 70mg/ day  
Narcolepsy 5 mg/day to 60 mg/day

### Methamphetamine dosing:

ADHD approved but not commonly used  
5 mg/day to 25 mg/ day

**Illicit use of amphetamines/  
methamphetamines up to 1 g / day**

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

105

105

## SOME CONSEQUENCES ARE DUE TO MODE OF CONSUMPTION

### » Smoking

- » Burned lips
- » Throat problems
- » Lung problems- acute (50% of those who smoke cocaine) and chronic 

### » Injection (unsafe practices)

- » Skin & heart infections
- » Hepatitis or HIV

### » Snorting

- » Sinus infections
- » Holes in nasal septum
- » Nosebleeds
- » Hoarseness

**NOTE:**  
There is cross tolerance from one class of stimulants to another

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

106

106

## EFFECTS DEPENDENT UPON MODE OF CONSUMPTION

### Drug Reaches Brain

- Smoking- seconds
- Injection- seconds
- Snorting- 15 minutes
- Oral-45 minutes

### Half-Life

- Cocaine 1h
- Bath Salts 3 hours
- Amphetamine 7 hours
- Methamphetamines 12 hours

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

107

107

## TIME FOR A POLL

Have you had trouble retaining patients with stimulant use disorders in treatment?

- A. Yes
- B. No

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

108

108

## STIMULANTS EFFECTS ON BRAIN CHEMISTRY


### Cocaine: Reuptake Blocker

INDIRECT agonist of

- + dopamine
- + norepinephrine
- + serotonin

### BLOCKS

- + neurotransmitters reuptake
- + sodium channels



### Amphetamines: Releaser

INDIRECT agonist of

- + dopamine
- + norepinephrine
- + serotonin

### INHIBITS

- + metabolism of neurotransmitters
- + vesicular storage
- + reverses reuptake

Photo Source: <https://www.drugabuse.gov/news-events/nida-notes/2017/03/impacts-drugs-neurotransmission>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

109

109

## ACUTE EFFECTS OF STIMULANTS

- Increased
  - Alertness/vigilance, concentration, mental acuity
  - Energy, locomotion
  - Sensory awareness & sexual desire
  - Self confidence, grandiosity, anxiety, irritability, paranoia
  - Heart rate & blood pressure, irregular heartbeat, vasoconstriction
  - Breathing rate, temperature, pupil size & blood sugar
  - Electrical activity, seizures
- Euphoria
- Abnormal bowel and bladder function

Toxic effects on muscles including  
Dystonia, tremors, stereotypy (i.e., ritualistic movements)

### Decreased

Brain blood flow & glucose metabolism  
Appetite & sleep  
Judgment & complex multi-tasking

### Cardiovascular effects

Heart attacks  
Arrhythmias  
Severe hypertension  
Strokes

Increased potential for violence and psychosis

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

110

110

## STIMULANT INTOXICATION: TREAT THE PRESENTING SIGN/SYMPTOM

### Overdose:

Seek immediate medical attention for:

- Hypertensive (HTN) crisis
- Cardiac arrhythmias
- Heart attack
- Stroke – Act F.A.S.T.\*
- Psychosis

### Treatment of Overdose

Treat HTN with alpha and/ or beta blockers

Treat arrhythmias with anti-arrhythmics

Treat vasoconstriction with nitroglycerin

### BH interventions for Overdose

Talk down the client in a calm environment

Treat agitation with benzodiazepine

Treat psychosis with antipsychotics

\* **F**acial drooping, **A**rm weakness, **S**peech difficulty, **T**ime to call 9-1-1

© 2023 Health Management Associates, Inc. All Rights Reserved.


HEALTH MANAGEMENT ASSOCIATES

111

111

## LONG-TERM MENTAL EFFECTS OF ILLICIT STIMULANTS

- » Tolerance to euphoria and appetite suppression
- » **Loss of ability to concentrate & severe memory loss**
- » Loss of ability to feel pleasure without drug



- » Paranoia and psychosis (hallucinations & delusions)
- » Insomnia and fatigue
- » Irritability and anger
- » **Depression (suicidal ideation)**
- » Impulsive, risky sexual behavior

\* Use of stimulants in doses approved by FDA for treatment of medical conditions do not result in these effects

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES


112

112



## LONG TERM PHYSICAL EFFECTS OF ILLICIT STIMULANTS

- » **Dry mouth, severe dental decay & gum problems**
- » **Bruxism (tooth grinding)**
- » Weight loss
- » Increased sweating; oily skin
- » Skin lesions from injection & formication (leading to skin picking)
- » Headaches
- » Movement disorders and Seizures
- » **Strokes (bleeding into the brain) & heart attacks**
- » Irregular heart beats
- » Cardiomyopathy
- » Kidney & liver failure
- » Pulmonary hypertension
- » Damaged brain cells
- » Neonatal effects



© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

113

113

## STIMULANTS AND PREGNANCY

- » Maternal death- pregnancy may increase risk of cardiovascular events
- » Preterm labor
- » Earlier gestational age at delivery
- » Low birth rate
- » Small for gestational age
- » Strokes in utero
- » Secreted in breast milk

**Child:**  
Dysregulated behavior, growth, inhibitory control, attention and abstract reasoning, but these effects appear to be related to gestational age at delivery, psychiatric disorders, other prenatal exposures and quality of postnatal environment. \*

**Anxiety, depression at 3-year-old \*\***  
**Worse cognitive function at 7-year-old \*\***

Source: Gouin 2011- cocaine; Kalaitzopoulos, 2018

\*Smid, M. C., Metz, T. D., & Gordon, A. J. (2019). Stimulant Use in Pregnancy: An Under-recognized Epidemic Among Pregnant Women. *Clinical obstetrics and gynecology*, 62(1), 168–184. <https://doi.org/10.1097/GRF.0000000000000418>

\*\*Deruf et al. 2007

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

114

114

## STIMULANT USE IN PREGNANT PEOPLE

### »» Pregnancy

- »» During pregnancy stimulant use is more common than opioid use
- »» Cannabis is the most used substance during pregnancy
  - »» Followed by stimulants

### »» Homelessness and sexual violence predict stimulant use in women...

If Post-traumatic Stress Disorder (PTSD) is present

- »» Integrated treatment is more effective for co-occurring disorder (COD)

#### Sources:

- Center for Behavioral Health Statistics Quality. 2015 National survey on drug use and health: Detailed tables In:2016
- Riley, ED. Risk factors for stimulant use among homeless and unstably housed adult women. Drug Alcohol Depend. 2015 August 1; 153: 173–179. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4510017/pdf/nihms694947.pdf>
- Ruglass LM, Hien DA, Hu M, Campbell ANC. Associations Between Post-traumatic Stress Symptoms, Stimulant Use and Treatment Outcomes: A Secondary Analysis of NIDA's Women and Trauma Study. Amer J on Addictions. Vol 23(1): 90-95. Jan-Feb 2014. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1521-0391.2013.12068.x>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

115

115

## CESSATION FROM STIMULANTS

- Acute withdrawal:
  - 4 days
  - No medication recommended
- Symptoms
  - Increased appetite
  - Increased sleep & dreaming
  - Decreased activity & energy
  - Depression & anhedonia
  - Decreased concentration
  - Craving

- Protracted withdrawal
  - Up to 10 weeks
  - No medication recommended
- Lingering effects on the brain; may be permanent
  - Psychosis
  - Movement Disorders
  - Cognitive Issues

Handout: Stimulant Withdrawal:  
Monitoring & Treatment  
<https://addictionfreeca.org/r/fpnseg8rpkgg>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

116

116

## AMPHETAMINES AND COGNITIVE IMPAIRMENT

- » Two-thirds of people with amphetamine use disorder have cognitive impairment
  - » Impairment is “associated” with:
    - » Older age
    - » Earlier onset of use
    - » Longer duration of use
    - » Greater frequency of use
  - » May limit ability to follow through on treatment
- » Damage cell structures
    - » Mitochondria in neurons & microglia
  - » Damage DNA
    - » Chromosomal alterations
  - » Inflammation of microglia
  - » Disruption of blood brain barrier
    - » Inflammatory markers in peripheral blood
  - » Cell death

Source: Paulus, M (2020) Neurobiology, clinical presentation, and treatment of methamphetamine use disorder a review. JAMA Psychiatry 77(9): 959-66.

117

## AMPHETAMINES AND LINGERING EFFECTS ON BRAIN

- » May be permanent even with prolonged abstinence
    - » Attention
    - » Memory
    - » Learning efficiency
    - » Visual- spatial processing
    - » Processing speed
    - » Psychomotor speed
    - » Executive dysfunction
- Cognitive Impairment**  
Impairs ability to engage in treatment due to trouble

  - Sequencing events to get to treatment
  - Remembering what is taught
  - Applying what is taught

118

## TREATMENT OF STIMULANT USE DISORDER

- » Harm Reduction needed due to IV use & risk of fentanyl
  - » Educational materials on psychological & physical effects
  - » Fentanyl test strips
  - » Syringe Exchange/distribution & other clean injection supplies
  - » Naloxone and overdose prevention education
  - » Quiet rooms to come down
  - » Showers & antibiotics for infection prevention & treatment
  - » Condoms & info on safe sex practices
  - » Water for hydration
  - » Toothpaste and toothbrush



Photo sources:  
Reproductive Health  
Supplies Coalition, Sara  
Groblechner, and Giorgio  
Trovato on Unsplash

119

119

## TREATMENT OF STIMULANT USE DISORDER: SAMHSA EVIDENCE BASED RESOURCE GUIDE

- » Motivational Interviewing (MI)
  - » Decreased days of stimulant use & amount of stimulant used/ day
- » Cognitive Behavior Therapy (CBT)
  - » Decreased quantity of stimulant use & frequency/ week
  - » Decreased risky sexual behaviors
- » Community Reinforcement Approach- see next slide
- » Contingency Management- see next slide

**STRONG EVIDENCE FOR THESE AS INDIVIDUAL INTERVENTIONS OR IN COMBINATION APPROACHES**

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

120

120

## TREATMENT OF STIMULANT USE DISORDER

- »» Community Reinforcement Approach (CRA)
  - »» Decreased addiction severity
  - »» Decreased drug use (weeks of use, frequency/week, \$/week)
  - »» Increased cocaine abstinence
- »» Contingency Management (CM): Strongest Effect Size
  - »» Decreased
    - »» days of stimulant use
    - »» stimulant cravings
    - »» HIV risk behaviors
  - »» Studies Veterans Administration National Rollout
    - »» Pre-CM: compared to 42% completed 2 sessions in 1 year
    - »» Post-CM Implementation: 50% completed 14 sessions in 12 week
    - »» 92% of >69,000 toxicology tests negative

Sources: SAMHSA  
Oliva, EM (2013)  
Warner & DePhilippis (2020)

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

121

121

## HOW DOES CM WORK?

**REMEMBER:**  
Measure objectively & frequently  
Don't set the bar too high or low

- »» Select objective target behavior (abstinence)
  - »» Define the behaviors
    - »» Attendance at clinic (group appt, urine)
    - »» Abstinence from DOC? all illicit drugs? prescribed drugs? alcohol?
- »» Provide immediate, consistent, tangible, desired rewards for target behavior
- »» Escalate size of reward for consistent behavior
- »» When target behavior does not occur
  - »» Withhold the reward
  - »» Reset size of reward for next occurrence of behavior
- »» Example: Fishbowl Method
  - »» 250 good job cards/gifts
  - »» 209 vouchers for \$1; 40 for \$20; 1 for \$100

**Reinforcement totaling  
\$80 = treatment as usual.  
Reinforcements of \$240  
improves outcomes.  
Petry 2004**

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

122


122

## IN THE CHAT BOX PLEASE ANSWER THIS QUESTION:

***Do you have a Contingency Management Program?***

**Yes**

**No**



© 2023 Health Management Associates, Inc. All Rights Reserved.

123

## WHAT TREATMENTS HAVE BEEN TRIED FOR STIMULANT USE DISORDER?

- » Replacement therapies: cocaine vs amphetamines
- » Antidepressants: SSRIs and tricyclic antidepressants not effective
- » Bupropion: risk of seizures; 5 failed trials for amphetamine use disorder \*
- » Mirtazapine: risk of weight gain; single small study + for amphetamine use disorder in men who have sex with men
- » Treatment of co-occurring Opioid Use Disorder (OUD)
- » Opioid agonists: increased dose of buprenorphine or methadone shows decreased cocaine use generally
- » Naltrexone: + results in multiple small studies amphetamine use disorder and cocaine use disorder \*
- » Antiseizure medications: Topiramate (risks); + one or two small studies in amphetamine use disorder

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

\* See next slide

124

124

## WHAT TREATMENTS HAVE BEEN TRIED FOR STIMULANT USE DISORDER?

[https://downloads.asam.org/sitefinity-production-blobs/docs/default-source/quality-science/stud\\_guideline\\_document\\_final.pdf?sfvrsn=71094b38\\_1](https://downloads.asam.org/sitefinity-production-blobs/docs/default-source/quality-science/stud_guideline_document_final.pdf?sfvrsn=71094b38_1)



The ASAM/AAAP  
**CLINICAL PRACTICE GUIDELINE ON THE**  
**Management of**  
**Stimulant Use**  
**Disorder**

**There are NO FDA approved medications for stimulant use disorders. Best Practices and Standards of Care do NOT endorse medication for stimulant disorders, by prescribers who are not experienced in addiction medicine.**

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

125

125

## WHAT'S ALL THE FUSS ABOUT?

- » New England Journal of Medicine article 2021
- » 400 adults with methamphetamine use disorder
- » Bupropion 450mg per day + placebo or bupropion 450mg per day + extended-release naltrexone 380mg IM q 3w (XR NTX )
- » 6-week study
- » Response defined as 3 of 4 toxicology tests negative for methamphetamines
- » 14% of patients on Bupropion + XR NTX responded vs 3% on Bupropion + placebo
- » Buprenorphine vs. placebo has a 21% difference for negative tox screen

**This is a 6-week study and has NOT been replicated yet. Only 11% increased response over placebo. Compare this to the EXCELLENT outcomes from psychosocial treatments.**

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

\* See next slide

126

126

## REFERENCES: STIMULANT USE

Baicy, K., & London, E. D. (2007). Corticolimbic dysregulation and chronic methamphetamine abuse. *Addiction*, 102, 5–15. doi:10.1111/j.1360-0443.2006.01777.x

Bolívar HA, Klempere EM, Coleman SRM, DeSarno M, Skelly JM, Higgins ST. Contingency Management for Patients Receiving Medication for Opioid Use Disorder: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. 2021 Oct 1;78(10):1092-1102. doi: 10.1001/jamapsychiatry.2021.1969. Erratum in: *JAMA Psychiatry*. 2022 Jan 26;: PMID: 34347030; PMCID: PMC8340014.

California Department of Public Health. (2020). California Opioid Overdose Surveillance Dashboard. <https://skylab.cdph.ca.gov/ODdash/>

CDC/NCHS, National Vital Statistics System, Mortality. CDC WONDER, Atlanta, GA: US Department of Health and Human Services, CDC; 2020. <https://wonder.cdc.gov/>

Chester, N., Mottram, D. R., Reilly, T., & Powell, M. (2004). Elimination of ephedrine in urine following multiple dosing: the consequences for athletes, in relation to doping control. *British journal of clinical pharmacology*, 57(1), 62–67.

DeFulio A, Rzeszutek MJ, Furgeson J, Ryan S, Reznia S. A smartphone-smartcard platform for contingency management in an inner-city substance use disorder outpatient program. *J Subst Abuse Treat*. 2021 Jan;120:108188. doi: 10.1016/j.jsat.2020.108188. Epub 2020 Nov 2. PMID: 33298295.

Foulds, J., & Young, J. T. (2019). Pharmacotherapy for incarcerated people with a history of violence: Response to commentary by Schofield et al. *Australian & New Zealand Journal of Psychiatry*, 54(1), 106–107. doi:10.1177/0004867419885175

Gouin, K., Murphy, K., & Shah, P. S. (2011). Effects of cocaine use during pregnancy on low birthweight and preterm birth: systematic review and metaanalyses. *American Journal of Obstetrics and Gynecology*, 204(4), 340.e1–340.e12. doi:10.1016/j.ajog.2010.11.013

Hedegaard H, Miniño AM, Warner M. Drug overdose deaths in the United States, 1999–2017. NCHS Data Brief, no 329. Hyattsville, MD: National Center for Health Statistics. 2018.

Hedegaard H, Miniño AM, Warner M. Drug overdose deaths in the United States, 1999–2018. NCHS Data Brief, no 356. Hyattsville, MD: National Center for Health Statistics. 2020.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

127

127

## REFERENCES: STIMULANT USE

Kalaiztopoulos, D.-R., Chatzistergiou, K., Amylidi, A.-L., Kokkinidis, D. G., & Goulis, D. G. (2018). Effect of Methamphetamine Hydrochloride on Pregnancy Outcome. *Journal of Addiction Medicine*, 12(3), 220–226. doi:10.1097/adm.0000000000000391

Lee, N. K., Jenner, L., Harney, A., & Cameron, J. (2018). Pharmacotherapy for amphetamine dependence: A systematic review. *Drug and Alcohol Dependence*. doi:10.1016/j.drugalcdep.2018.06.038

Mayo Clinic Laboratories. (2020). Amphetamine-Type Stimulants (ATS). <https://www.mayocliniclabs.com/test-info/drug-book/amphetamine.html>

NCHS, National Vital Statistics System. Estimates for 2019 and 2020 are based on provisional data. Estimates for 2015-2018 are based on final data (available from: [https://www.cdc.gov/nchs/nvss/mortality\\_public\\_use\\_data.htm](https://www.cdc.gov/nchs/nvss/mortality_public_use_data.htm))

NIDA. 2020, April 3. California: Opioid-Involved Deaths and Related Harms. Retrieved from <https://www.drugabuse.gov/drug-topics/opioids/opioid-summaries-by-state/california-opioid-involved-deaths-related-harms> on 2020, October 7

NIHCM Foundation. (2020). Beyond Opioids: Rapid Increase in Drug Deaths Involving Stimulants. <https://www.nihcm.org/categories/beyond-opioids-rapid-increase-in-drug-deaths-involving-stimulants>

Oliva, EM, Bowe, T., Harris, A. H., & Trafton, J. A. (2013). Datapoints: False starts in psychotherapy for substance use disorders and PTSD in the VHA. *Psychiatric Services*, 64(8), 722.

Paulus, M (2020) Neurobiology, clinical presentation, and treatment of methamphetamine use disorder a review. *JAMA Psychiatry* 77(9): 959-66.

Peirce, JM, et al. (2006). Effects of lower-cost incentives on stimulant abstinence in methadone maintenance treatment: A National Drug Abuse Treatment Clinical Trials Network study. *Archives of General Psychiatry*, 63(2), 201–208. <https://doi.org/10.1001/archpsyc.63.2.201>

Petry N., T. J. (2004). Prize reinforcement contingency management for treating cocaine users: how low can we go, and with whom? *Addiction*, 349-360.

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

128

128



## REFERENCES: STIMULANT USE

Substance Abuse and Mental Health Services Administration. (2017). Medical Review Officer Guidance Manual for Federal Workplace Drug Testing Programs. [https://www.samhsa.gov/sites/default/files/workplace/mro-guidance-manual-oct2017\\_2.pdf](https://www.samhsa.gov/sites/default/files/workplace/mro-guidance-manual-oct2017_2.pdf)

Substance Abuse and Mental Health Services Administration (SAMHSA). Treatment of Stimulant Use Disorders. SAMHSA Publication No. PEP20-06-01-001 Rockville, MD: National Mental Health and Substance Use Policy Laboratory. Substance Abuse and Mental Health Services Administration, 2020. <https://store.samhsa.gov/product/Treatment-of-Stimulant-Use-Disorder/PEP20-06-01-001>

Sever, P. S., Dring, L. G., & Williams, R. T. (1975). The metabolism of (?)-ephedrine in man. *European Journal of Clinical Pharmacology*, 9(2-3), 193–198. doi:10.1007/bf00614017

Smid, M. C., Metz, T. D., & Gordon, A. J. (2019). Stimulant Use in Pregnancy: An Under-recognized Epidemic Among Pregnant Women. *Clinical obstetrics and gynecology*, 62(1), 168–184. <https://doi.org/10.1097/GRF.0000000000000418>

U.S. Drug Enforcement Administration, Diversion Control Division. (2019). NFLIS-Drug Special Report: Methamphetamine Reported in NFLIS, 2001–2017. Springfield, VA: U.S. Drug Enforcement Administration. <https://www.nflis.deadiversion.usdoj.gov/DesktopModules/ReportDownloads/Reports/12568NFLISdrugMethamphetamine.pdf>

Volkow, N. D., & Morales, M. (2015). The Brain on Drugs: From Reward to Addiction. *Cell*, 162(4), 712–725. doi:10.1016/j.cell.2015.07.046

Werner, D. D. and DePhilippis, D. (2020, August 12). Contingency Management for People Experiencing Homelessness - Homeless and Housing Resource Network SAMHSA Webinar. Washington DC, Washington DC.

Disorder stimulant use management of - downloads.asam.org. (n.d.). [https://downloads.asam.org/sitefinity-production-blobs/docs/default-source/quality-science/stud\\_guideline\\_document\\_final.pdf?sfvrsn=71094b38\\_1](https://downloads.asam.org/sitefinity-production-blobs/docs/default-source/quality-science/stud_guideline_document_final.pdf?sfvrsn=71094b38_1)

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

129

129

## CHEMSEX

130

130

## SUD AND HIV RISK

- » The co-occurrence of HIV and SUD in a community increases the risk of HIV transmission due to:
  - » Sharing of syringes
  - » Intoxicant and/or stimulant involved unprotected sex
  - » Sexual violence and victimization
  - » Unaware of HIV status
  - » Unsuppressed viral load

*HIV can be a risk factor for substance use.  
But also...  
Substance use can be a risk factor for HIV transmission.*

131

## CHEMSEX

### Definition:

Chemsex (also known as sexualized drug use – SDU) is the **use of drugs to enhance sexual experience**. Common drugs used include methamphetamine, gamma-hydroxybutyrate (GHB), gamma-butyrolactone (GBL), cocaine, ketamine, poppers (amyl nitrite) or cannabis (the latter two gave rise to the term SDU)

### What You Should Know:

- Chemsex is popular among some gay, bisexual, transgender, and queer persons, **but can be experienced by persons of any gender**
- Chemsex participants have higher odds of condomless anal sex with partners of different or unknown HIV status (bareback sex)
- Persons engaged in Chemsex have greater risk of acquiring sexually transmitted infections (STIs) and hepatitis C (HCV)
- Participants are at higher risk of HIV transmission
- The association with sexual risk indicates the importance of promoting harm reduction among this population (e.g., condoms, PrEP, PEP, drug knowledge).

132

## CHEMSEX

### Common Terminology Used to Communicate the Desire to Engage in Chemsex

Injecting	Meth	GHB	Ketamine	ChemSex
Pointing, slamming, darts	Blowing clouds, Cloudy, ice cream, tea, T, tina	Water, Gina, Swirling	K, Special K	Party, PNP, Party and play
	  	 		  

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

133

133

## METHAMPHETAMINE AND ITS IMPACT ON HIV INFECTION

Methamphetamine use:

- » Increases sexual desire, impairs judgment, and provides energy and confidence to engage in sexual activity for long periods of time (hyper-sexual)
- » Causes erectile dysfunction
- » Causes mucosal dryness
- » Decreases adherence to HIV treatment and medical follow-up
- » Increases HIV replication
- » Accelerates progress of HIV-related dementia

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

134

134

## DOES METHAMPHETAMINE ACCELERATE HIV AND HCV?

- » In test tube studies, when methamphetamine is added to immune cells, it significantly increases HIV replication
  - » Particularly in CD4 cells and monocytes (white blood cells)
- » In mouse models, methamphetamine activated a portion of the HIV genetic code (long terminal repeat – LTR), prompting cells to release a protein tied to more rapid HIV disease progression
- » The Journal of Viral Hepatitis published a study indicating that methamphetamine increases Hepatitis C replication.

Source: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2675873/>

135

## HIV AND HEPATITIS C


136

## HIV AND HEPATITIS C CO-INFECTIONS

**Hepatitis C and HIV**  
are often-overlooked consequences of America's **opioid crisis**.

**EIGHT IN TEN**

**new Hepatitis C infections** in the U.S. are transmitted through **injection drug use**.



Nearly **ONE IN TEN**


**new HIV infections** in 2015 were due to **injection drug use**.

HepVu.org      SOURCE: U.S. CENTERS FOR DISEASE CONTROL AND PREVENTION      HepVu

© 2023 Health Management Associates, Inc. All Rights Reserved.      HEALTH MANAGEMENT ASSOCIATES      137

137

## HIV AND HEPATITIS C CO-INFECTIONS



**Nine states** represent **more than half, or**

**52%**

**of all people living with Hepatitis C nationwide.**

Estimated Number of People Living with Hepatitis C, 2013-2016

0 - 10,000	10,001 - 25,000	25,001 - 50,000	50,001 - 75,000	75,001+
------------	-----------------	-----------------	-----------------	---------

HepVu.org      SOURCE: HepVu      HepVu

© 2023 Health Management Associates, Inc. All Rights Reserved.      HEALTH MANAGEMENT ASSOCIATES      138

138

## HIV AND HEPATITIS C CO-INFECTIONS

- » In 2018 in Minnesota, there were 60 acute HCV cases and 33,856 chronic cases
  - » 8,140 Co-infected for HIV and HCV
- » The U.S. Public Health Service/Infectious Diseases Society of America guidelines recommend that all HIV-infected persons be screened for HCV infection (CDC, 2014).

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

139

139

## QUESTIONS?

140

140

## NEXT STEPS

- » Join us for Session 4 next Wednesday!
- » Your registration should have included a reoccurring calendar invite for all four sessions
- » Please complete the evaluation and post-test for this session that will be sent out after via email (evaluations must be completed for those seeking CEU/CME credits).

Follow-up questions?

Contact Carlos Mena at [cmena@healthmanagement.com](mailto:cmena@healthmanagement.com)

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

141

141

## AGENDA FOR WEBINAR SERIES

SESSION	TOPICS
<b>#1</b> <b>WEDNESDAY, NOV 1</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> Understanding HIV <input type="checkbox"/> HIV Testing and Treatment <input type="checkbox"/> The Science of Addiction <input type="checkbox"/> Screening, and Assessment
<b>#2</b> <b>WEDNESDAY, NOV 8</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> Ethical and Legal Issues <input type="checkbox"/> Funding and Policy Considerations <input type="checkbox"/> HIV Risk Reduction <input type="checkbox"/> SUD Harm Reduction <input type="checkbox"/> HIV and Stigma <input type="checkbox"/> Motivational Interviewing
<b>#3</b> <b>WEDNESDAY, NOV 15</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> Working with Justice Involved Persons <input type="checkbox"/> Substance Use Disorder Treatment with Medications <input type="checkbox"/> Mental Health Treatment and Counseling <input type="checkbox"/> Stimulant Use <input type="checkbox"/> Chem Sex
<b>#4</b> <b>WEDNESDAY, NOV 22</b> 12:00 pm to 3:00 pm	<input type="checkbox"/> <b>Cultural, Racial and Sexual Identities</b> <input type="checkbox"/> <b>Pregnancy and HIV, SUD/ODU</b> <input type="checkbox"/> <b>Accessing, Obtaining, and Integrating Services for Individuals with HIV and SUD in Minnesota</b>

© 2023 Health Management Associates, Inc. All Rights Reserved.

HEALTH MANAGEMENT ASSOCIATES

142

142